



REGIONAL SEAS

GESAMP:

***Cadmium, lead and tin
in the marine environment***

UNEP Regional Seas Reports and Studies No. 56

GESAMP Reports and Studies No. 22

Prepared in co-operation with



United Nations



FAO



UNESCO



WHO



WMO



IMO



IAEA

PREFACE

GESAMP, the Joint Group of Experts on the Scientific Aspects of Marine Pollution, was established in 1969 and is today co-sponsored by the International Maritime Organization (IMO), Food and Agriculture Organization of the United Nations (FAO), United Nations Educational, Scientific and Cultural Organization (UNESCO), World Meteorological Organization (WMO), World Health Organization (WHO), International Atomic Energy Agency (IAEA), United Nations and United Nations Environment Programme (UNEP). According to its present terms of reference, the functions of GESAMP are:

- to provide advice relating to the scientific aspects of marine pollution ^{1/}; and
- to prepare periodic reviews of the state of the Marine environment as regards marine pollution and to identify problem areas requiring special attention.

Since its beginning GESAMP involved a large number of experts as members of GESAMP or GESAMP Working Groups and produced, at the request of the sponsoring organizations, numerous reports ^{2/}.

This document reproduces the substantive part of the report of the GESAMP Working Group on Review of Harmful Substances, approved by the fourteenth session of GESAMP (Vienna, 26 - 30 March 1984).

The Working Group was chaired consecutively by Messrs B.H. Ketchum, A. Jernelöv, and L. Friberg. The following experts participated in the preparation of the report: Dr. B.G. Bennett, Prof. L. Friberg, Prof. A. Furtado Rahde, Prof. A. Jernelöv, Mr. R. Lloyd, Dr. L. Magos, Prof. S.P. Meyers, Dr. A. Oskarsson, Dr. P.M. Sivalingam, Prof. G. Tomassi, Dr. H. Galal-Gorchev, Dr. M. Gilbert, Dr. R. Helmer, Dr. J. Parizek, and Dr. G. Vettorazzi.

The Working Group was requested

- to prepare short and referenced reviews on selected substances which include an assessment of the following factors:
 - (a) the total amount of the particular substance(s) which reach(es) the marine environment (on a local, regional, and global scale) with particular attention being given to the relative importance of land-based sources;

^{1/} GESAMP defined marine pollution as "introduction by man, directly or indirectly, of substances or energy into the marine environment (including estuaries) resulting in such deleterious effects as harm to living resources, hazards to human health, hindrance to marine activities including fishing, impairment of quality for use of sea-water, and reduction of amenities."

^{2/} V. Pravdic: GESAMP, The First Dozen Years. UNEP, 1981.

- (b) the fate (transport, distribution, and transformation) of the substance in the marine environment; and
 - (c) the effects of the substance on the marine environment and adjacent coastal areas, including direct and indirect effects on living resources, human health and amenities;
- to produce a scientific evaluation of the harmful effects of substances released into the marine environment on living resources, human health, aesthetics, and other legitimate uses of the marine environment and adjacent coastal areas.

The activities of the Working Group were organized by WHO, acting as the "lead agency". The Working Group was jointly sponsored by WHO, FAO, and UNEP.

CONTENTS

	<u>Page</u>
I. INTRODUCTION	1
1. Evaluation mechanisms	1
2. Working procedures of the group	1
3. Quality of data base	1
3.1 Analytical quality control	1
3.2 Ecotoxicological quality aspect	2
3.3 Quality of human toxicological data base	2
4. Dietary intake considerations	2
4.1 Basis for total dietary intake estimates	2
4.2 Seafood consumption patterns	3
5. References	4
II. CADMIUM	6
1. Cadmium in the marine environment	6
1.1 Reference documentation	6
1.2 General facts	6
1.3 Sources	6
1.4 Transport, transformation, and bioaccumulation	9
1.4.1 Transport	9
1.4.2 Transformation	9
1.4.3 Bioaccumulation	10
1.5 Cadmium in sea water, sediments, and marine biota	12
1.5.1 Sea water	12
1.5.2 Sediments	12
1.5.3 Marine biota	12
2. Effects on marine biota	14
2.1 Reference documentation	14
2.2 Effects on marine biota	15
3. Human health aspects	17
3.1 Reference documentation	17
3.2 Toxicokinetic properties	17
3.3 Health effects	19
3.4 Total exposure to cadmium	20
3.5 Contribution of cadmium from marine food	20
3.6 Evaluation of potential health effects	21
4. Conclusions on cadmium	22
4.1 Potential harm to living resources	22
4.2 Potential hazards to human health	22
5. References	23
III. LEAD	31
1. Lead in the marine environment	31
1.1 Reference documentation	31
1.2 General facts	31
1.3 Sources	31
1.4 Transport, transformation, and bioaccumulation	33
1.4.1 Transport	33
1.4.2 Transformation	34
1.4.3 Bioaccumulation	34
1.4.3.1 Inorganic lead	35
1.4.3.2 Organic lead compounds	36
1.5 Lead in sea water, sediments, and marine biota	37
1.5.1 Sea water	37
1.5.2 Sediments	37

	<u>Page</u>	
2.2	Effects on marine biota	40
2.2.1	Inorganic lead	40
2.2.2	Organic lead	42
3.	Human health aspects	42
3.1	Reference documentation	42
3.2	Toxicokinetic properties	43
3.3	Health effects	46
3.4	Total exposure to lead	48
3.4.1	Dietary lead intake	49
3.4.2	Exposure from water and air	51
3.5	Contribution of lead from marine food	51
3.6	Evaluation of potential health effects	52
4.	Conclusions on lead	52
4.1	Potential harm to living resources	52
4.2	Potential hazards to human health	53
5.	References	53
IV.	TIN	63
1.	Tin in the marine environment	63
1.1	Reference documentation	63
1.2	General facts	63
1.3	Sources	63
1.4	Transport, transformation, and bioaccumulation	64
1.4.1	Transport	64
1.4.2	Transformation	64
1.4.3	Bioaccumulation	66
1.5	Tin and organotin concentrations in sea water, sediments, and marine biota	69
1.5.1	Sea water	69
1.5.2	Sediments	69
1.5.3	Marine biota	70
2.	Effects on marine biota	70
2.1	Reference documentation	70
2.2	Effects on marine biota	70
2.2.1	Inorganic tin	70
2.2.2	Organic tin	71
3.	Human health aspects	73
3.1	Reference documentation	73
3.2	Toxicokinetic properties	73
3.2.1	Inorganic tin	73
3.2.2	Organotin compounds	74
3.3	Health effects	74
3.3.1	Inorganic tin	74
3.3.2	Organotin compounds	74
3.4	Total exposure to tin	75
3.5	Contribution of tin from marine food	75
3.6	Evaluation of potential health effects	76
4.	Conclusions on tin	76
4.1	Potential harm to living resources	76
4.2	Potential hazards to human health	77
5.	References	78
ANNEX I	83

I. INTRODUCTION

1. Evaluation Mechanisms

From an examination of data profiles by UNEP's International Register of Potentially Toxic Chemicals (IRPTC), other data profiles, and available critical reviews of published data, significant papers were selected by the Working Group for thorough evaluation. These papers, together with recent and pertinent publications, then formed the basis of this review. It is recognized, however, that these papers provide only a partial coverage of the world literature. Information was lacking in several areas essential to an environmental hazard evaluation of cadmium, lead, and zinc, and these areas were identified in the review.

2. Working Procedures of the Group

The method and approaches applied by the Working Group were discussed and agreed upon at a planning session in Stockholm, 24-25 September, 1982. This was attended by the chairmen of GESAMP and of the Working Group, and by international agency representatives.

For each substance, selected experts prepared draft sections of the review. The reviews were then critically examined and revised by the Working Group members (Annex I). The final draft was submitted to GESAMP for consideration, comments, and adoption.

3. Quality of Data Base

3.1 Analytical quality control

Many studies conducted in the various countries aimed at evaluating normal and elevated levels of trace metals in different media. Unfortunately, most published reports lack quality assurance data, and valid comparisons cannot, therefore, be made. Furthermore, results from several inter-laboratory comparisons amplify the need for quality control. A review of such comparison studies, with particular emphasis on lead and cadmium in blood, has recently been published in connection with a UNEP/WHO Biological Monitoring Project on Assessment of Human Exposure to Lead and Cadmium through Biological Monitoring (Vahter, 1982). Various intercalibration exercises with those laboratories engaged in the determination of trace metals in commercially important marine organisms from the North Atlantic were also organized, since 1971, by ICES, the International Council for Exploration of the Sea (Topping, 1983). These reviews clearly show that errors may be large even in "experienced" laboratories. Since 1975, the IAEA's International Laboratory of Marine Radioactivity has also operated a large global analytical quality control programme for metals and chlorinated hydrocarbons in marine organisms and sediments. A similar programme has been run by the Intergovernmental Oceanographic Commission for Seawater Samples.

The introduction of sophisticated and increasingly sensitive analytical techniques has made it possible to measure trace substances in extremely low concentrations. Simultaneously, however, the risks of interference from competing factors has increased considerably. Although the awareness of the need for quality control has also increased during recent years, it is not possible to state, generally, that analyses carried out during the last 5-year period, for example, are always more reliable than earlier analyses.

Analytical problems may occur with any matrix to be examined. Particular problems arise, however, when analysing biological media or matrices which have very low trace metal concentrations (picograms/g). Schaule & Patterson (1980) showed that, for example, the lead concentrations in seawater samples have been overestimated by factors of up to 5000 and, that the lead concentrations reported for marine organisms are, with very few exceptions, several orders of magnitude higher than the actual concentrations present. These high concentrations are caused by the sample becoming contaminated during analysis. The importance of implementing rigid quality assurance programmes was amplified in 2 recent trace metal programmes sponsored by UNEP/WHO. One measured lead and cadmium in blood, and cadmium in kidneys (Vahter, 1982; Friberg & Vahter, 1983).

first programme, it was rare that a laboratory met the criteria for data acceptance throughout the training phase, and gross errors were often recorded. The food study noted that the results of current analytical quality control analyses allowed few conclusions to be drawn concerning the reliability of previously collected data. In addition to the various forms of analytical error, there is the possibility of contaminating biological samples, for example, by use of unsuitable sample collection vials and contaminated chemicals. Furthermore, errors due to adsorption and desorption on the walls of containers may cause inaccurate results.

The various sources of error which are possible make it necessary to exercise great caution when evaluating analytical data. In particular, it is more the exception than the rule that data on quality assurance are presented as part of published studies. Such caution has been exercised in the evaluation carried out by the Working Group, but there is still no guarantee that all the data used in this evaluation are completely valid. If rigid quality assurance criteria had been required, the analytical data available for use in the evaluation would have been extremely limited.

3.2 Ecotoxicological quality aspect

Experiments on marine organisms were carried out using many different procedures and techniques, and the usefulness of the results to the present review was critically examined. In particular, only a few experiments offered analytical confirmation of the concentrations and various forms of a substance that the subject was exposed to. There was little data on proven harmful effects resulting from chronic exposure. Therefore, extrapolations from the limited data base in order to predict whether environmental effects are likely to occur have to be treated with caution. In this respect, there is also a need for reliable analytical data on environmental concentrations for some species of the substances evaluated.

3.3 Quality of human toxicological data base

The quality and quantity of toxicological data show substantial variation from one marine pollutant to another. Ideally, the evaluation of the health hazard presented by a certain pollutant ought to be based on data which include a comprehensive dose-effect relationship. For a selected and preferably critical effect, a reliable dose-response curve should be provided. Equally important is an established correlation between the concentration of the toxic chemical (or one of its metabolites) in an index medium and the effects and responses. Some examples relevant to the substances selected by GESAMP are presented in the following paragraphs.

For one of the most widely studied metals, lead, the relationship between blood lead and the effect of lead on the synthesis of haemoglobin is well established, but the relationship between blood lead and the effect of lead at the lower end of exposure on the development of the Central Nervous System (CNS) in children is a question of controversy. The relationship between oral lead intake and blood lead concentration has not been investigated, and the lack of this information hinders the prediction of blood lead concentration from daily dietary intake and vice versa.

The corresponding correlation between blood cadmium and current exposure has been established. In a condition of changing exposure, however, blood cadmium may not correlate with the renal concentration of cadmium. However, the quality of data that predicts kidney accumulation of cadmium from dietary intake has recently become more reliable.

In many cases, the evaluation is dependent on experimental animal data, especially in experiments which aim to study those quantitative relationships which can be extrapolated to man. Unfortunately, in the cases of tin and particularly of organotins, even the animal data base is not sufficient for extrapolation, and dose-related human data are totally absent.

4. Dietary Intake Considerations

4.1 Basis for total dietary intake estimates

Accurate estimates of dietary intake of food contaminants are difficult to make for

contaminant levels and rates of consumption. It is possible, however, to estimate intake from measurements of the concentration of contaminants in specific foods and the amounts of food consumed. Alternatively, measurements may be made with composite samples of the total diet. This approach requires fewer measurements and may reflect actual intake more closely, but it has the disadvantage of disregarding the contribution of single foods to the total dietary intake of contaminants. The calculated dietary intake may be made for representative population exposures, or it may be made with special reference to critical groups, for example, children, pregnant women, high consumers, etc. From contributions of particular food items to the total dietary intake, it is useful to note the "critical" foods to which more attention should be given in surveillance programmes.

Comparisons between calculated dietary intake and tolerable intake limits may indicate safety levels, or the incidence of risk for the exposed population. Due to the large variations in food consumption patterns among countries and to the wide variations in food contaminant concentrations, mean dietary intakes should be calculated and evaluated at the national or even local level.

Finally, the total energy value and composition of the diet should be taken into account in an evaluation of potential risks for population groups. It has been demonstrated that not only total food intake but also components such as fat, calcium, iron, and zinc can influence a subject's susceptibility to toxicity of contaminants by modifying the degree of gastrointestinal absorption.

4.2 Seafood consumption patterns

Seafoods do not represent a significant component of the diet for much of the world's population. Marine foods consumed by man include many trophic levels. Seaweeds are eaten mainly in the Far East, but also in Europe, for example, as laverbread or agar. Phytoplankton and zooplankton are not themselves consumed, but their predators, such as oysters, mussels, clams, scallops, herring, and sardines are. However, krill is used for animal fodder. Higher trophic levels include the carnivorous gastropods, clams, cephalopods, crabs, and shrimp, and foremost, fish. The world (population 4150 million in 1977) average of fish and shellfish protein consumption is 3.8 g per person per day of a total of 68.8 g total protein per person per day (FAO, 1980a). These data can also be converted to and analysed in terms of the fresh weight of marine foods. The world's average daily consumption of 3.8 g protein corresponds to about 20 g edible fish and shellfish per day or 140 g edible tissue, which is equal to about one meal of fish and shellfish, per week. Aquatic plants and seaweeds make up only 4% of the world's harvest of marine and freshwater foods. Of the total catches of aquatic animals, 10% are caught in fresh water and the rest is marine. Seventy-seven percent of the world's landings are marine fish, 7% are molluscs, and 4% are crustaceans, with other marine animals contributing 1% or less. The northwest Pacific and northeast Atlantic are the most productive areas (FAO, 1980b).

From the Table on seafood consumption in selected countries (Table 1), it can be seen that the small populations on the islands mentioned eat relatively large amounts of marine foods. According to FAO's food balance sheets (1980a), the population (40 000 inhabitants) of the Faeroe Islands consumes, on the average, 38.6 g protein (equal to 193 g fresh weight of edible tissue) originating from marine foods. Assuming that 150 g of edible tissue constitutes one meal, this corresponds to more than one meal per day of seafood. The average Japanese intake is the highest among larger populations. By comparison, the average consumption of marine foods in Australia, the USA, and the USSR is small. However, certain individuals are reported to consume much greater amounts. Fisherman at sea, especially in less affluent regions, will consume exclusively marine foods. For example, about 800 g per day were consumed on fishing boats in southern Italy (Bernhard & Renzoni, 1977). Canadian Indians are believed to have an intake of up to 1300 g per day during the fishing season, while 800 - 1500 g per day were eaten by fishermen and their families at Minemata (Review: Piotrowski & Inskip, 1981). Average consumption levels have been estimated on a global basis to be 23 g per day per person, with 38 g per day as the value for Europe.

Attention must be paid to situations in which seafood consumption represents a

Table 1. Consumption of fish and seafood (g fresh weight) (living weight) and in percent of different types of marine food^a

	Population	Total marine protein	Seafood in fresh weight	% of different types of marine foods on protein basis		
Australia	13.8	3.4	17	68	24	
Bermuda	0.06	15.8	79	87	13	
Faeroe Islands	0.04	38.6	193	82	16	3
Iceland	0.1	19.2	96	83	13	
Japan	113.9	22.5	113	77	4	10
Maldives	0.14	37.1	186	100		
Portugal	8.8	10.2	51	87	2	3
USA	216.8	3.2	16	63	22	6
USSR	258.9	9.4	47	81		1
Vanuatu	0.1	23.2	116	99	2	
Yemen Republic	1.9	13.0	65	100		

^a From: Bernhard & Andreae (1983).

Note: The protein values (FAO, 1980a) have been converted into fresh weight, assuming that 100 g fresh weight contain 20 g protein.

example, be the case with mercury in certain species of fish, or that with cadmium in oysters. It has been calculated, for example, that, while the contribution of seafoods to the total dietary intake of cadmium and lead is only 6% for the general population in Italy, the percentage can increase up to 25% for cadmium and 40% for lead in fishermen living in the Italian coastal villages where fish consumption is about 10 times that of the average quantity of seafood consumed in the rest of Italy.

5. References

BERNHARD, M. & ANDREAE, M.O. (1983) Transport of trace metals in marine food chains. In: Nriagu, J.O., ed. Changing metal cycles and human health, Berlin, Springer Verlag, pp. 143-167.

BERNHARD, M. & RENZONI, A. (1977) Mercury concentration in Mediterranean marine organisms and their environment: natural or anthropogenic origin. Thalassia Jugosl., 13: 265-300.

FAO (1980a) Food balance sheets and per caput food supplies, Rome, Food and Agriculture Organization of the United Nations.

FAO (1980b) Yearbook of fishery statistical catches and landings, Rome, Food and Agriculture Organization of the United Nations, Vol. 48.

FRIBERG, L.T. & VAHTER, M. (1983) Assessment of exposure to lead and cadmium through biological monitoring results of a UNEP/WHO global study. Environ. Res., 30: 95-128.

GESAMP (1976) Review of harmful substances, New York, United Nations, 80 pp (Rep. Stud. GESAMP No. 2).

NATIONAL FOOD ADMINISTRATION (1982) Summary and assessment of data received from the FAO/WHO Collaborating Centres for food contamination monitoring, Uppsala, Sweden, National Food Administration.

PATTERSON, C.C. & SEATTLE, D.M. (1976) The reduction of orders of magnitude errors in lead analysis of biological materials and natural waters by evaluating and controlling the extent and sources of industrial lead contamination introduced during sample

PIOTROWSKI, J.K. & INSKIP, M.J. (1981) Health effects of methylmercury, London, Chelsea College (MARC Report No. 24).

SCHAULE, B. & PATTERSON, C.C. (1980) The occurrence of lead in the northeast Pacific and the effects of anthropogenic inputs. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 345-352.

VAHTER, M., ed. (1982) Assessment of human exposure to lead and cadmium through biological monitoring, Stockholm, Liber Tryck (Prepared for United Nations Environment Programme and World Health Organization by National Swedish Institute of Environmental Medicine and Department of Environmental Hygiene, The Karolinska Institute, Stockholm).

II. CADMIUM

1. Cadmium in the Marine Environment

1.1 Reference documentation

The major reviews and reference works used were Aylett (1973) on the chemistry of cadmium, Frei & Hutzinger (1976) on analytical techniques, Webb (1979) on general chemistry and biology, WHO (1979) on environmental health criteria, Friberg et al. (1974) and Fleischer et al. (1974) on cadmium in the environment and its impact, and Simpson (1981) for a critical review of cadmium in the marine environment. On bioaccumulation, reviews by Alabaster (1978) and Coombs (1979) were consulted. The kinetics of uptake were reviewed by McLeese (1980), and George (1980) reviewed the pertinent literature on mussels.

Many other individual papers by research workers were consulted and are listed in the reference section.

1.2 General facts

Cadmium (Cd) (Greek Cadmean (earth), calamine) is in subgroup IIb, Zn, Cd, Hg of the transition series in the Periodic Table of Elements. It has atomic number 48 and atomic weight 112.40. Cadmium was first isolated and identified by F. Strohmeyer in 1817 from the zinc ore smithsonite ($ZnCO_3$). It has been released into the environment since the early days from the smelting of a variety of ores and the burning of wood and coal. Cadmium is among the rarer trace elements and is seldom found in pure minerals. It is extracted commercially from zinc ores, e.g., zincblende (ZnS), in which it occurs at 0.1 - 5.0%.

Cadmium is more mobile in non-polluted, undisturbed soils than, for example, lead. It is even more accessible and more mobile in cultivated soils under the many influences of soil chemistry (Page et al., 1981).

The predominant state of oxidation in nature is Cd^{2+} , which is a borderline Type (b) cation. In freshwater, cadmium is extensively associated with colloidal and particulate matter, and soluble speciation is confined to the free Cd^{2+} ion together with small amounts of $CdCl_2$ and $CdSO_4$. In the sea, some 66% of cadmium is present as free Cd^{2+} together with $CdCO_3$ (26%), $Cd(OH)_2$ (5%), $CdCl_2$ (1%), and $CdSO_4$ (1%) (Whitfield et al., 1981). In coastal and estuarine waters, a high proportion of cadmium is associated with particles and is present as complexes (Nriagu, 1980; MacKay, 1983).

1.3 Sources

Typical cadmium concentrations found in igneous rocks are $0.001 - 1.8 \mu g g^{-1}$ (mean, $0.15 \mu g g^{-1}$), in metamorphic rocks $0.04 - 1.0 \mu g g^{-1}$, in sedimentary rocks, $0.3 - 11 \mu g g^{-1}$, in shales, up to $90 \mu g g^{-1}$, in marine clays, $0.4 \mu g g^{-1}$, and in marine phosphorites, $60 - 340 \mu g g^{-1}$ (Page et al., 1981; Simpson, 1981). Agricultural soils from unpolluted areas usually contain less than $1 \mu g g^{-1}$. These data, derived from sundry original sources, reflect the extreme values that were found and may possibly include analytical discrepancies. Areas in which enhanced levels of cadmium are found are usually linked with the occurrence of zinc-rich ore bodies, zinc smelting, and other zinc-related manufacturing processes and metal plating operations. Localized and naturally high cadmium concentrations may be found near deposits of sulfide ores such as sphalerite, phosphorite, hydrothermally-mineralized rocks, and some black shale deposits such as in the United Kingdom and California, USA.

Smaller but important sources of cadmium are a by-product of copper refining and, to a lesser extent, lead processing. Natural emissions of cadmium to the atmosphere are associated with volcanic eruptions, such as that of Mount Etna (Buat-Menard & Arnold, 1978), and forest fires and windblown dusts (Simpson, 1981; Hutton, 1982). Similarly, cadmium is released into the atmosphere by power generation facilities which use fossil

Cadmium in water comes from contaminated agricultural soils, mining wastes, mine waters, and the industrial use of cadmium. An important source is municipal sewage effluents and sludges, including those of domestic origin.

World production of cadmium metal was about 18 900 tonnes in 1979. The main producers are the USA, the USSR, Japan, and Canada, followed by Belgium and France.

The main anthropogenic sources relate to ore mines, metallurgical industries, and to the disposal of sewage sludges. Cadmium concentrations in the fumes of copper, lead, nickel, and zinc sulfide smelters can be relatively high due to the high volatility of the metal (Fleischer et al., 1974). Other major atmospheric inputs come from the combustion of fossil fuels in industries using coke and from the incineration of domestic refuse (Fleischer et al., 1974), as shown in Table 2. These atmospheric inputs have demonstrated effects on agricultural soils and products in the surrounding areas (Peterson & Alloway, 1979). In addition, cadmium may be introduced by urban and motorway dusts, cadmium-contaminated phosphorus-containing fertilizers, and sewage sludge applications on land.

Table 2. Summary of current cadmium inputs to the environment of the European Commission Countries (tonnes year⁻¹)^a

Source	Compartment		
	Air	Land	Water
Volcanic action	20	ND	ND
Non-ferrous metal production			
zinc and cadmium	20	200	50
copper	6	15	ND
lead	7	40	20
Production of cadmium-containing materials	3	90	108
Iron and steel production	34	349	ND
Fuel combustion			
coal and lignite	8	390	ND
oil and gas	0.5	14.5	-
Waste disposal	31	1434	ND
Sewage sludge disposal	2	130	33
Phosphate fertilizers	-	346	62
Totals	132	3009	273

^a From: Hutton (1982).

Cadmium is a scarce and fairly expensive metal of low mechanical strength. It is released slowly into the environment from widespread sources. Cadmium is mainly applied via electroplating or dipping to another metal as a thin film coating for protection against corrosion. It is also much used as a pigment in yellow or brown paints and cadmium metal is used in special alloys and solders. Seldom is it possible to recover the metal economically. Use of cadmium in alkaline Ni-Cd rechargeable batteries has potential environmental hazards in view of the amounts of nickel and cadmium involved in

Table 3. World consumption of cadmium by main uses, 1965, 1970, and 1975^a

Use	1965		1970		1975	
	tonnes	%	tonnes	%	tonnes	%
Batteries	669	7	842	8	1102	14
Pigments	2463	25	2733	25	1980	25
Stabilizers	905	9	2089	19	1249	16
Plating	4518	47	4068	37	2614	33
Alloys	804	8	803	7	658	8
Others	333	4	382	4	300	4
Total	9692		10 917		7903	

^a Sources: S.A. Hiscock, Cadmium Association, 34 Berkeley Square, London.

There is a growing application of organocadmium compounds in the plastics industry. Alkyl-cadmiums are used as polymerizing catalysts in pvc manufacture. Cadmium laurate, stearate, palmitate, myristate, and others are used to reduce weathering effects on plastics. The rate of release of organocadmium complexes to the aquatic environment from these sources is likely to be low. However, the simpler methyl-cadmium (Me₂Cd) is decomposed rapidly in air and water.

The reasons why sewage (domestic and mixed) may contain high proportions of cadmium relative to other trace metals are not clear, nor is the reason why the cadmium content varies irregularly (Peterson & Alloway, 1979). The cadmium content of 189 samples from 150 wastewater treatment plants in the USA ranged from 3 to 3410 $\mu\text{g g}^{-1}$ dry weight sludge, with a strong positive correlation to the degree of industrialization in the area observed. The mean value was 16 $\mu\text{g g}^{-1}$ (Sommers, 1977). Förstner & van Lierde (1979) list values for Sweden, England, and Wales, and Michigan, USA in the lower part of this range, and the more recent ranges for England (Murray et al., 1980) are lower than the above average. The US Food and Drug Administration recommends, along with other restraints, an upper limit of 29 $\mu\text{g g}^{-1}$ for sewage sludge applied to agricultural land.

Relatively high levels of cadmium are found in dredged spoils. Data from dredging in US waters (Krenkel et al., 1976) range from 0.6 to 4.1 $\mu\text{g g}^{-1}$ dry weight (8 sites) with 17.6 $\mu\text{g g}^{-1}$ near Long Island Sound. Ranges were found in the Clyde and other Scottish estuaries for various harbour silts of 5.6 - 6.8 $\mu\text{g g}^{-1}$. Murray & Norton (1979) report many values in the range 0.2 - 8.2 $\mu\text{g g}^{-1}$. Using their data, they calculated an annual input of 17 tonnes cadmium for 1977 in the United Kingdom, on the basis that a total of 28×10^6 tonnes soil (14×10^6 tonnes dry solids) is dumped into the North Sea and Irish Sea from harbour and channel dredging in the country.

Studies on the transfer of the different forms of cadmium from freshwater to the sea are inconclusive. Certain estuaries are conservative in most trace metals (Bewers & Yeats, 1981). The daily inputs of cadmium to inshore waters of the North Sea from the Humber estuary (Table 4) have been assessed by Murray et al. (1980). The relative proportions and magnitudes of these inputs should be of general interest to other fairly heavily urbanized and industrialized estuary areas.

The estimated inputs to the environment of the European Community (Table 2) give a firm indication of emissions to the air and disposals to land but only an incomplete account of aquatic inputs. The generalized global anthropogenic input of cadmium to the

Table 4. Comparison of cadmium inputs to the Humber Estuary^a

Rivers	15 kg day ⁻¹	40.4%
Sewage discharges	2 kg day ⁻¹	5.4%
Industrial discharges	8 kg day ⁻¹	21.5%
Sewage sludge dumping	0.3 kg day ⁻¹	0.8%
Industrial waste dumping	0	0%
Dredged spoil	6.8 kg day ⁻¹	18.3%
Atmospheric input	5 kg day ⁻¹	13.5%
Direct coastal discharges	0.05 kg day ⁻¹	0.1%
Total	37.15	100%

^a Extract from Table 5 of Murray et al. (1980).

1.4 Transport, transformation, and bioaccumulation

1.4.1 Transport

Cadmium enters the seas and oceans from the air mainly in particulate form and, to a lesser extent, dissolved in rain and snow. Wittmann (1979) quotes the enrichment of cadmium in atmospheric particulate matter relative to the earth's crust as 300 in north Atlantic westerly winds and 1900 in the urban air of the USA. In some remote areas, volcanic emissions may be the principal source of enrichment. The concentration of cadmium in samples of air, above the Atlantic Ocean between Iceland and the Bermudas, ranged from 0.003 to 0.62 ng m⁻³ (Duce et al., 1976). This concentration range is comparable with that of samples collected in rural areas of the USA (Fassett, 1980). Much higher levels of cadmium in air are observed in urban and industrialized areas, particularly near metal refining and processing plants. Airborne cadmium is a principal source of input to offshore and oceanic waters.

The transport of cadmium from freshwater to the sea occurs either in particulate or soluble form. The specific form depends on the state of the river, its mineralization and its sources of pollution, as well as on unidentified local factors. Quantification is difficult. The general range for cadmium content in clean rivers and lakes is about 0.1 - 1.2 µg litre⁻¹; in polluted industrial rivers in the United Kingdom, the USA, and in Europe, it is 1 - 36 µg litre⁻¹ (Coombs, 1979). Freshwaters respond to natural or anthropogenic exposure to metalliferous ores, soils, and sediments. Values of 1 - 9 µg litre⁻¹ were found for the Jintsu river in Japan which flows through the area where the Itai-itai disease occurred; 30 µg litre⁻¹ was found in the drainage streams of a nearby ore mine. In the United Kingdom, 3 - 95 µg litre⁻¹ was found in streams in North Wales, while 5 - 20 µg litre⁻¹ was found in areas in Cornwall affected by ore mining. Rivers are subjected to affected by cadmium inputs from raw and treated sewage, for example, the River Rhine has a range of 1 - 10 µg litre⁻¹ (Coombs, 1979).

River sediments generally reflect the neighbouring soils and mineral workings. High cadmium levels are invariably accompanied by high levels of other trace metals. As a result of these inputs, there are enhanced levels of cadmium in near-shore sediments and sea waters (see below).

1.4.2 Transformation

The transformation of inorganic forms of cadmium in sea water scarcely affects its

of the element, but this has not been demonstrated. Organic chelates, such as humates, are likely to liberate bound cadmium as the result of dilution and degradation in sea water. Cadmium is held in sewage sludge partly in combination with carbonates and sulfides, and partly in complex organic combinations. In the latter example, cadmium combines with the sulfur-rich fractions of organic matter of which there is great excess, for the inorganic and organic contributions are widely variable (Stover et al., 1976; Sommers et al., 1977). Since most biogenic cadmium-organic complexes, including metallothioneins, are fairly easily biodegradable (Coombs, personal communication, 1983), there is a ready release of cadmium into aerobic waters and sediments; under anaerobic conditions, insoluble cadmium carbonates and sulphides may persist in sediments.

1.4.3 Bioaccumulation

Simpson (1981) regards the processes of uptake (or absorption) by phytoplankton, followed with grazing by herbivores and the subsequent elimination of cadmium in faecal pellets, as a major contributory factor affecting cadmium distribution in the photic zone. Such a mechanism might also be the cause of elevated concentrations (up to 60 $\mu\text{g g}^{-1}$) of cadmium in superficial sediments of the Walvis Bay, and might possibly explain the unusually high concentration, up to 600 $\mu\text{g g}^{-1}$, for Red Sea sedimentary deposits in an anaerobic environment created by a massive bloom.

Many estimates are available for bioaccumulation in marine flora and fauna (Coombs, 1979). For many species, covering most phyla, accumulation factors are of the order of thousands. For some molluscs and some arthropods, they are tens of thousands, and for certain tissues (of which few, if any, are usually eaten by man), they are hundreds of thousands.

Aquatic organisms can be exposed to cadmium in the ambient water, in sediments, and in their diet. The bioavailability of cadmium and, consequently, its accumulation in tissues depends on a number of factors. Non-complexed cadmium added to a rapidly growing algal culture has slightly less effect on the growth rate than when added at the time of inoculation of the medium (Kayser & Sperling, 1980). In experiments with the American oyster, complexed cadmium was found to be less readily accumulated (Yen-Wan Hung, 1982); the lower concentration used, however, 40 - 60 $\mu\text{g cadmium litre}^{-1}$, may also have contributed to this effect.

Ray et al. (1980) found that the rate of cadmium uptake by the polychaete Nereis virens, when exposed to contaminated sediments (1 - 4 mg cadmium kg^{-1}), was the same as its uptake from solutions that contained the same cadmium concentrations (30 - 100 $\mu\text{g cadmium litre}^{-1}$) as that present in the water overlying the sediments. Further studies by Ray et al. (1981b) showed that little accumulation from cadmium-contaminated sediments occurred in Nereis virens, Macoma balthica, and Crangon septemspinosa. Similar studies showed that N. virens, Mercenaria mercenaria, and Palaemonetes pugio did not accumulate cadmium from contaminated sediments during a 100-day exposure period (Rubinstein et al., 1983). Hardy et al. (1981) exposed excised gills of the clam Protothaca staminea to contaminated sediments and interstitial water and concluded that "low level additions of cadmium to sea water are not likely to lead to significant bioaccumulation through the gills of suspension-feeding bivalves". Therefore, cadmium bound in contaminated sediments does not appear to be bioavailable to marine organisms. However, molluscan herbivores may accumulate cadmium from littoral algae, and this may be accumulated, in turn, by carnivores feeding on these organisms (Davies, 1981; Simpson, 1981).

Algae

Phytoplankton can accumulate significant concentrations of cadmium (Kremling et al., 1978; Kayser & Sperling, 1980), although decomposing cells rapidly release cadmium into the water. It is assumed that the metal is loosely bound to the cell's surface, and this process appears to be important in the biogeochemical cycling of cadmium in the marine system. Laminaria saccharina, in common with other seaweeds, can also accumulate significant concentrations (Markham et al., 1980).

Crustaceans

Shrimps and lobsters do not appear to accumulate cadmium from ambient aqueous concentrations of less than about $2 \mu\text{g litre}^{-1}$ (data reviewed in McLeese, 1980), although White & Rainbow (1982) found some evidence of accumulation at lower concentrations in artificial seawater. The hepatopancreas is a major site of accumulation of cadmium from polluted waters containing higher concentrations (Ray et al., 1981a); studies by Davies (1981) have shown that accumulation at this site is derived from cadmium-contaminated food and not from cadmium in solution. Depuration rates in clean water range from a half-life of 11 days to no loss (data reviewed in McLeese, 1980). Von Bias (1981) found that the amphipod Corophium volutator accumulated cadmium more readily at low salinities, and that no more than 50% was lost on return to clean water.

Molluscs

Greatest attention has been given to this group of organisms, especially bivalves, because of their linear rate of cadmium uptake and their tolerance to high body burdens which makes them good sentinel species. Linear uptake rates have been recorded by von Westernhagen et al. (1978) in Mytilus edulis exposed to $5 \mu\text{g cadmium litre}^{-1}$ for 163 days; exposure of this species to 10 and $100 \mu\text{g cadmium litre}^{-1}$ for 17 days also resulted in linear uptake. Body burdens of up to 150 mg kg^{-1} did not affect respiration or growth rates (Poulson et al., 1982). Uptake rates increased with temperature in Crassostrea virginica (Zarogian, 1980) and Saccostrea echinata (Denton & Burden-Jones, 1981). These 2 authors also recorded increased uptake at lower salinities. Uptake rates can depend on feeding intensity which may be reduced slightly at high cadmium concentrations (Ward, 1982) and are related to ventilation rates as well (Janssen & Scholz, 1979); intermittent emersion of Mytilus edulis appears to increase the rate of uptake during the immersed period as compared with those which are continuously immersed (Coleman, 1980).

Cadmium appears to be bound to the metallothionein in mollusc tissues. In the kidney tissue of Bay scallop (Argopecton irradians) exposed for 5 days to $700 \mu\text{g cadmium litre}^{-1}$, the concretions (or granules) contained 60% of the accumulated cadmium, and only 2% was bound to low molecular weight proteins (Carmichael & Fowler, 1981). In addition, in Mytilus edulis which was exposed to $100 \mu\text{g cadmium litre}^{-1}$ for 3 months, George & Pirie (1979) found that 85% of the cadmium in membrane-limited granular structures may have been associated with metallothionein. In the same species, Marshall & Talbot (1979) found cadmium associated with sulfur and sometimes phosphorus in membrane-bound vesicles.

It is evident that cadmium accumulated from low environmental concentrations can be rapidly bound into a non-toxic complex (Carpene & George, 1981) which is retained within the body and excreted very slowly, if at all. Mowdy (1981) found that 50% of accumulated cadmium in Crassostrea virginica was lost in 60 days in clean water (the rate of loss being slower at low salinities), and George & Coombs (1977) showed that the cadmium excretion rate in Mytilus edulis was 18 times slower than the uptake rate. Denton & Burden-Jones (1981), also, found that the cadmium half-life in Saccostrea echinata was very long. Mussels loaded with $564 \text{ mg cadmium kg}^{-1}$ dry weight lost 47 mg kg^{-1} in a 42-day depuration period, in which time the fraction bound to metallothionein rose from 22 to 78% (Kohler & Riisgard, 1982). However, because accumulation occurs in tissues which may be sites of toxic action, harmful effects may follow when the cadmium binding sites are saturated.

Fish

Cadmium accumulates in the liver and gills of plaice (Pleuronectes platessa) but not in their muscle; liver concentrations began to increase only after 70 days exposure to $5 \mu\text{g cadmium litre}^{-1}$ (von Westernhagen et al., 1978). Similar data were obtained for plaice by Pentreath (1977) and for dabs (Limanda limanda) by von Westernhagen et al. (1980). Few data are available for depuration rates of cadmium from fish. However, Noel-Lambot (1981) found that the gut of several species of fish contained "intestinal corpuscles" - a mixture of mucous cells, mucous, and granules - which have a high

Pentreath (1977) found that 4 days after feeding plaice with cadmium-loaded Nereis, about 5% of the ingested cadmium was associated with the gut wall and none detected in other internal organs. After defaecation, the half-life of the remaining cadmium was between 100 and 200 days.

Birds and mammals

Elevated concentrations of cadmium have been found in the liver and kidney of sea birds (Bull et al., 1977), seals, and porpoises (Falconer et al., 1983), but in no instances have these levels been identified as causing harmful effects. The levels are thought to result from a natural accumulation through the food chain.

1.5 Cadmium in sea water, sediments, and marine biota

1.5.1 Sea water

Many observations of the cadmium content of sea water in estuaries, inshore waters, seas with restricted circulation, shelf waters, and open oceans are detailed in the review by Simpson (1981). In general, the concentration in sea water is about 0.01 - 0.1 $\mu\text{g cadmium litre}^{-1}$, (Preston et al., 1972; Campbell & Loring, 1980; Magnusson & Westerlund, 1980; Simpson, 1981). Values which are about 5 - 10 times higher, 0.2 - 0.4 $\mu\text{g cadmium litre}^{-1}$, have been reported from certain coastal areas, such as the Oslofjorden, Norway, and Liverpool Bay, United Kingdom (Preston et al., 1972; Rojahn, 1972). The very local nature of estuarine variability was already described, and the broad pattern of declining land influences can be traced in decreasing cadmium concentrations towards the open ocean (Table 5).

In the more stable water regime of the open ocean a significant reduction in cadmium, parallel to the reduction in phosphate and nitrate, can be demonstrated in the upper water layer which is influenced by the photic zone and phytoplankton productivity (Simpson, 1981).

1.5.2 Sediments

The same progression is even more evident in the cadmium content of bottom sediments (Nicholson & Moore, 1981; Simpson, 1981). Extremely high values are reported for enclosed waters affected by local ore mining (US east coast, before dredging, 30 - 18 400 $\mu\text{g g}^{-1}$; after dredging, 40 - 5000 $\mu\text{g g}^{-1}$), but values in fjords, harbours, and estuaries with marked industrial and urban influence are more commonly in the range 10 - 1000 $\mu\text{g g}^{-1}$. In many of the cleaner inshore areas, a fairly broad range (about 0.2 - 5.0 $\mu\text{g g}^{-1}$) is common at each site, with upper values extending to 10 $\mu\text{g g}^{-1}$ where mineralization and other influences, such as oil activities, are present. Sediments in the Baltic Sea range from 0.2 - 2.2 $\mu\text{g g}^{-1}$, while the values for the North Sea are below 1.0 $\mu\text{g g}^{-1}$ for 80% of the samples (Nicholson & Moore, 1981), with indications of higher values near sites of oil-related activity.

In the open oceans, most values are below 0.5 $\mu\text{g g}^{-1}$; Aston et al. (1972) report a mean of 0.23 $\mu\text{g g}^{-1}$ for cadmium in deep-sea sediments from the North Atlantic and a higher mean of 0.65 $\mu\text{g g}^{-1}$ for samples taken near the mid-Atlantic Ridge. An important feature to take into account is the influence of upwelling. Diatomaceous ooze from Walvis Bay, a major upwelling area, contained 3 - 60 $\mu\text{g g}^{-1}$. This compared with samples farther offshore of diatomaceous ooze which showed 0.17 - 0.88 $\mu\text{g g}^{-1}$ and of radiolarian ooze which had 0.13 - 0.98 $\mu\text{g g}^{-1}$. An extensive Atlantis II survey of the deeps in the Red Sea also yielded high values: 2 - 600 $\mu\text{g g}^{-1}$ and 30 - 3900 $\mu\text{g litre}^{-1}$ in the interstitial water (Simpson, 1981).

1.5.3 Marine biota

Phytoplankton

It is particularly difficult to determine accurately the cadmium content of the phytoplankton which are the first stage in the marine food web. Likely values do not appear to exceed a few $\mu\text{g g}^{-1}$ on a dry weight basis (Kremling et al., 1978; Kavser &

Table 5. A summary of cadmium distribution in saline waters^a

	Background	"Normal" range (dissolved, $\mu\text{g litre}^{-1}$)	Suspected contamination	Top of range	Background	"Normal" range (sediment, $\mu\text{g g}^{-1}$)	Suspected contamination	Top of range
Estuaries and closed bays	0.01	0.05 - 0.2	> 0.2	45.7	< 1.0	0.1 - 2	> 2.0	50 000
Bays and coastal waters	< 0.01	0.01 - 0.15	> 0.15	10.3 (100)	< 1.0	0.1 - 1.5	> 1.5	60
Open sea	< 0.01	0.01 - 0.1	> 0.1	1.6 0.65	< 1.0	0.1 - 1.0	> 1.0	(600)
Open ocean	< 0.005	0.01 - 0.1	> 0.1	(1.61)	< 1.0	0.1 - 1.0	> 1.0	0.977

^a From: Simpson (1981).

g^{-1} , but brown algae contain a wide range ($0.2 - 26 \mu g g^{-1}$), which is believed to reflect the ambient cadmium concentration.

Zooplankton

The cadmium content of protozoa, parazoa, cnidaria, euphausiids, and chaetognaths is normally well below $2 \mu g g^{-1}$, although a single sample of copepods (crustacea) had a content of $9.8 \mu g g^{-1}$. Arthropods such as the sea-skaters (Halobates sp) have shown a notably high cadmium content with values of $50 - 210 \mu g g^{-1}$ (Bull et al., 1977) and $1.7 - 120 \mu g g^{-1}$ (Schulz-Baldes & Cheng, 1980). These insects live on and feed in the upper microlayer of the seas or oceans where the atmospheric cadmium input may have a special influence.

Molluscs

Among the molluscs, the edible gastropods such as limpets (Patella sp), ormers (Haliotis sp), and whelks (Buccinum sp), which live in the intertidal and subtidal zones, can contain exceptionally high levels of cadmium, namely $0.2 - 295 \mu g g^{-1}$ for limpets and ormers, and $0.05 - 730 \mu g g^{-1}$ for whelks. The bivalves, shellfish which are commercially important, can also contain the following moderately high cadmium concentrations: mussels (Mytilus edulis) and scallops (Pecten maximus) have values of $0.04 - 140 \mu g g^{-1}$, cockles (Cardium edule and other species), oysters, and clams show $0.3 - 170 \mu g g^{-1}$. Other important commercial shellfish (cephalopods), cuttle-fish, squids, and octopuses can have exceptionally high liver cadmium contents, $23 - 1100 \mu g g^{-1}$, but the decapods (shrimps, prawns, lobsters and, particularly, crabs) have a lower range, $0.5 - 3.3 \mu g g^{-1}$. The cadmium value of the brown meat of Cancer pagurus has values up to $9.1 \mu g g^{-1}$ wet weight in certain areas (Davies, 1981). Most concentrations of cadmium in molluscs vary greatly with location. In fact, mussels are widely used in monitoring and are useful indicators of pollution. The Joint Monitoring Programme of European Countries reports general results for Mytilus edulis of less than $0.7 \mu g g^{-1}$ wet weight. Values up to $15 \mu g g^{-1}$ have, however, been reported in localized regions (Paris Commission, 1983).

Fish

Regular monitoring has ascertained that most commercial fish species contain very low concentrations of cadmium (Murray, 1979; Davies, 1981). A recent comprehensive study from the Swedish National Food Administration, accompanied by a quality assurance programme, showed mean cadmium levels ranging from 1 to $27 \mu g$ cadmium kg^{-1} . There are reports of somewhat higher levels (Coombs, 1979), but there is no information related to quality assurance. The results of the Joint Monitoring Programme of European Countries for 1978-80 has shown generally very low concentrations of cadmium in muscle of fish, independent of species. The values were mostly less than $20 \mu g kg^{-1}$ wet weight (Paris Commission, 1983).

Mammals and birds

Cadmium levels are not markedly high in the top predators such as the common porpoise (Phocoena phocoena). Mean values found are, for male liver, 0.15, for male kidney, 1.1, for female liver, 0.27, and for female kidney, $2.7 \mu g g^{-1}$ wet weight (Falconer et al., 1983). Value ranges for the grey seal (Halichoerus grypus) are, for liver, $0.07 - 8.5 \mu g g^{-1}$ wet weight and, for kidney, $0.10 - 15 \mu g g^{-1}$ wet weight (McKie et al., 1980). Somewhat higher cadmium levels are reported for Atlantic seabirds (fulmar, Manx shearwater, puffin, Leach's petrel, storm petrel, razorbill). For example, in the liver were found $1.4 - 57 \mu g g^{-1}$ dry weight and, in the kidney, $15 - 240 \mu g g^{-1}$ dry weight were found, allowing for wet/dry weight basis (Bull et al., 1977).

2. Effects on Marine Biota

2.1 Reference documentation

Information consulted included the reviews by Alabaster (1978) and the US EPA (1980) and the tabulations of data on concentrations and effects summarized by Taylor (1981)

2.2 Effects on marine biota

Data given in the summaries by Taylor (1981) and IRPTC (1981) indicate that cadmium is not very toxic within short exposure periods, and the 96-h LC₅₀s for a wide range of species are usually in excess of 1 mg cadmium litre⁻¹. Similarly, chronic effects usually become apparent at concentrations greater than 50 µg cadmium litre⁻¹. It is most unlikely that these concentrations will occur in the future, even in the most polluted situations.

However, some species have been reported to be affected at cadmium concentrations less than 15 µg litre⁻¹ and usually after prolonged exposure under laboratory conditions. It is these data which form the basis of this critical review.

Algae

The growth rate of the dinoflagellate Prorocentrum micans was inhibited by 1.2 µg cadmium litre⁻¹ with resulting cell numbers in the cultures being less than one tenth of the control values; no effect was found at 0.4 µg cadmium litre⁻¹ (Kayser & Sperling, 1980). However, Prévot (1980) found that the growth rate of this species was only slightly affected at 5 µg cadmium litre⁻¹ and then only after 22 days exposure. A 50% reduction occurred at 60 µg litre⁻¹ with 30 days exposure. The reasons for this difference in results are not known. Concentrations greater than 10 µg cadmium litre⁻¹ were found to increase the vacuolation and number of lysosomes in this species (Soyer & Prévot, 1981). Li (1980) recorded a reduced growth rate of Isochrysis galbana when exposed to 1 µg cadmium litre⁻¹ for 10 days. Kayser (1982) found that 10 µg cadmium litre⁻¹ temporarily reduced the growth rate of Scrippsiella faeroense, but there was no such effect at 2.0 µg litre⁻¹. In contrast, Fisher & Froud (1980) found that 25 µg cadmium litre⁻¹ had no effect on the growth of 4 other species of diatoms, and Kremling et al. (1978) found that 1 µg cadmium litre⁻¹ had no effect on phytoplankton communities in a mesocosm. Other data indicate that concentrations greater than 100 µg cadmium litre⁻¹ are required to produce effects on a wide range of algal species.

Coelenterates

At a salinity of 10 ‰ and at 17.5 °C, irreversible retraction of 50% of hydranths of Laomedea loveni occurred at about 3 µg cadmium litre⁻¹, whereas the same effect was produced at 15 µg litre⁻¹ at 25 ‰ salinity (Theede et al., 1979). Other hydrozoa tested appear to be more resistant to cadmium.

Growth and survival of the ctenophore Pleurobrachia pileus were adversely affected by 1 µg cadmium litre⁻¹ in a marine mesocosm (Kuiper, 1981).

Annelida

This group of organisms appears to be very resistant to cadmium, no effects being recorded at concentrations less than 100 µg litre⁻¹ (Kuiper, 1981).

Crustacea

Embryonic development of the mud crab Eurypassopeus depressus to the megalopa stage was not affected by 10 µg cadmium litre⁻¹, but subsequent development to the crab stage was delayed and increased mortalities were noted. Phototactic swimming rates at Stages II and III were higher in larvae exposed to cadmium (Mirkes et al., 1978). Copepod populations grew in a mesocosm when exposed to 5 µg cadmium litre⁻¹ due to a decrease in predation by ctenophores; but exposure to 50 µg cadmium litre⁻¹ reduced their numbers (Kuiper, 1981).

The estuarine mysid (Mysidopsis bahia) appears to be very sensitive to cadmium with a 17-day LC₅₀ of 11 µg litre⁻¹, and a 50% reduction in numbers of young produced per female occurred in 6.4 µg litre⁻¹, although there was no effect at 4.8 µg litre⁻¹ (Nimmo et al., 1978). Further studies by Gentile et al. (1982) on this

Experiments with the lobster Homarus americanus showed that malate dehydrogenase activity intensified with exposure to 6 μg cadmium litre⁻¹ for 30 days; the MDH:LDH ratio was also higher (Gould, 1980). Similar experiments showed that isolated gill respiration rates increased when lobsters were exposed to 3 μg cadmium litre⁻¹ for 30 days, but no effects on osmoregulation were found (Thurberg et al., 1977). Using the brown shrimp Crangon crangon, Price & Uglow (1980) found an increase in scaphognathite rate during a 13-day exposure to 5 μg litre⁻¹. This is said to be close to the incipient lethal level, although no experimental data are given.

Molluscs

Although molluscs are resistant to cadmium toxicity, with effects on adults and larval stages being recorded at concentrations usually greater than 70 μg litre⁻¹, Zarogian & Morrison (1981) found that 5 μg litre⁻¹ slightly delayed the development of 10% of Crassostrea virginica larvae. However, after 3 weeks of exposure, the growth rates in both 5 and 15 μg cadmium litre⁻¹ solutions were similar to those of the controls. Similarly, Watling (1982) found that 10 μg cadmium litre⁻¹ caused a reduction in growth rate of C. margarita, which increased to above the control rate when returned to clean water. Much higher concentrations were required to produce this effect in C. cucullata and C. gigas. Stromgren (1982), too, found that 10 μg cadmium litre⁻¹ reduced the growth rate of mussels (Mytilus edulis) within a 9-day exposure period; 5 μg litre⁻¹ had no significant effect, and 2 μg litre⁻¹ stimulated growth to 145% of the control value.

Echinoderms

This group is resistant to cadmium. No harmful effects were recorded at concentrations less than 100 μg cadmium litre⁻¹; this included genotoxicity (Pagano et al., 1982).

Tunicates

Recent experiments show that Botryllus schlosseri is very resistant to cadmium (Kayser, 1982). Sub-lethal effects occurred only after prolonged exposure to more than 5 mg cadmium litre⁻¹.

Fish

Ojaveer et al. (1980) found that eggs of Baltic herring began to hatch earlier when exposed to 5 μg cadmium litre⁻¹, and newly-hatched larvae were 10% smaller than the controls; the significance of these small differences is not clear. Plaice (Pleuronectes platessa) exposed for 280 days to 5 μg cadmium litre⁻¹ grew more slowly than the controls between the 70- and 130-day exposure period; the feed ration in both groups was reduced during this period. After 130 days, the growth rate of all fish were similar (von Westernhagen et al., 1978). Hyperactivity of plaice was noted among the cadmium-exposed fish; a similar effect was noted in flounders (Platichthys flesus) that were exposed to 500 μg cadmium litre⁻¹ but not those exposed to 50 μg cadmium litre⁻¹ (Larsson et al., 1981). Exposure of plaice to 50 μg cadmium litre⁻¹ resulted in a 90% mortality rate within 96 days and an LT₅₀ of 30 days. Mortality among dabs (Limanda limanda) similarly exposed was 30% in 96 days, but no effects on growth of the survivors were noted (von Westernhagen et al., 1980). Mortality was thought to be caused by secondary infection of fin erosions. Weis & Weis (1976) noted that 10 μg cadmium litre⁻¹ temporarily inhibited caudal fin regeneration in the killfish Fundulus heteroclitus.

Larsson (1975) found that the blood haematocrit and haemoglobin content of flounders was unaffected in a 4-week exposure to 5 μg cadmium litre⁻¹, but a 26% reduction from control values occurred within the next 5 weeks. On the other hand, Calabrese et al. (1975) found that 5 and 10 μg cadmium litre⁻¹ had no effect on the haematological parameters of the winter flounder Pseudopleuronectes americanus within 60 days, although the oxygen consumption rate of excised gills was 15% lower than the control values. In similar experiments with this species, Gould (1977) found that

potassium content of blood plasma in flounders exposed to 5 µg cadmium litre⁻¹ for 9 weeks, but increases in phosphorus and magnesium took place only at 50 µg cadmium litre⁻¹. There was no effect on bone structure. Oxygen consumption of excised gills of striped bass (*Morone saxatilis*) exposed to 5 µg cadmium litre⁻¹ for 30 days was 27% lower than the control values, although this effect was neither apparent at 90 days nor was it apparent in gills from fish exposed to 2.5 µg cadmium litre⁻¹. No effect on the enzymes AAT and G6PDH was noted during the exposure period, although there was a decrease in their concentration when fish exposed to 5 µg cadmium litre⁻¹ were returned to clean water for 30 days (Dawson et al., 1977). It is not clear whether any of these changes can be considered as harmful.

3. Human Health Aspects

3.1 Reference documentation

The major reviews on cadmium consulted in the preparation of this section include Friberg et al. (1974), Tsuchiya (1978), Simpson (1981), Friberg et al. (in press), and the interim report on Environmental Health Criteria for Cadmium (WHO, 1979).

3.2 Toxicokinetic properties

The average absorption of cadmium from food is approximately 5% (Rahola et al., 1972), but people suffering from anaemia or those on a calcium-deficient diet may have a considerably higher rate of absorption (up to 20%) (Flanagan et al., 1978).

Cadmium is transported via the blood to other parts of the body. In blood, cadmium is mainly found in the red cells, where it is bound to a protein of low molecular weight, metallothionein. Most cadmium is stored in liver and kidneys where, after long-term low-level exposure, approximately 50% of the cadmium is to be found. About one third of the body burden is located in the kidneys, where the concentration in kidney cortex is probably about 1.25 times the average concentration in the whole kidney (Kjellström et al., 1984). Accumulation also takes place in certain other tissues, such as the muscle (Kjellström, 1979), where the biological half-time is long; concentrations are usually low, but with large amounts of tissue, an important part of the body burden is accounted for.

The placenta serves as an effective barrier against cadmium uptake, and the newborn is practically free from cadmium. The total body burden at birth is only about 1 µg, but continuous accumulation takes place in the body up to about the age of 50. At this age, the total body burden is between 10 and 30 mg with concentrations in the kidney cortex of 15 - 50 mg cadmium kg⁻¹. In countries where exposure via food is high, such as Japan, the total body burden and concentrations in kidney cortex may be considerably higher.

The biological half-time in the kidneys is long, approximately 20 years and possibly somewhat shorter at old age, which explains the accumulation of cadmium. In blood, part of the cadmium is related to body burden and has a long biological half-time. As a rule, the major part of cadmium in blood is related to recent exposure and has a half-time of about 2 - 3 months. The concentration in blood is, therefore, a useful indicator of exposure during recent months. The concentration in blood is low, usually below 1 µg cadmium litre⁻¹ in non-smokers, but may reach several µg cadmium litre⁻¹ in smokers. If exposure remains constant, blood levels may also be used for evaluation of long-term risks (WHO, 1980).

Due to the long biological half-time of cadmium in the body, only a small part of cadmium attributable to long-term low-level exposure will be excreted. Excretion takes place via faeces and urine and comprises only 0.005 - 0.1% per day of the total body burden. Since only a minor portion of ingested cadmium is absorbed, the amount excreted via faeces can be used to estimate total daily intake. Concentrations in urine will increase with age. As for long-term low-level exposure, excretion is related to body burden and concentrations in the kidney cortex. Cadmium levels in urine are not usually good indicators of recent exposure. When concentrations in kidney cortex reach critical levels and signs of kidney dysfunction occur, the excretion of cadmium increases dramatically. Concomitantly, the cadmium concentration in the kidneys decreases. This




Fig. 1. Concentration of cadmium in kidney cortex (geometric mean values with 1.28 times the geometric standard deviations indicated) in relation to smoking habits among the subjects (30 - 69 years of age) studied in Belgium, India (data from Ahmedabad, Bangalore, and Calcutta pooled), Japan, Yugoslavia. Swedish data from Elinder et al. (1976). Number of smokers (including former smokers) and non-smokers as well as mean age of subjects in each subgroup are indicated under the bars.

concentrations in the kidneys which are very low. At high exposure levels, such as can be seen in workers of certain industries, metabolism is different. Relatively more cadmium is accumulated in the liver, and concentrations in the blood and urine may be more difficult to interpret as indicators of exposure and body burden. For a detailed discussion of the metabolic model for cadmium, reference is made to Kjellström & Nordberg (1978) and Camner et al. (1979).

Very limited data are available on the different forms of chemical binding of cadmium to proteins or other compounds in foodstuffs and the influence this may have on the toxicokinetics of cadmium. Recent studies show cadmium is bound to different proteins in different species of oysters. In New Zealand Bluff oysters, cadmium is bound to a protein similar to metallothionein. People with extreme consumption of such oysters, leading to a daily cadmium intake of 200 - 500 µg cadmium, were found to have disproportionately low blood cadmium levels (McKenzie et al., 1982). This may indicate that the distribution of cadmium bound to metallothionein is different from unbound cadmium. Animal data support this view. Cadmium metallothionein administered parenterally or perorally is transported directly to the kidneys (Cherian & Shaikh, 1975; Nordberg et al., 1975; Cherian et al., 1978). Thus, blood cadmium may not reflect the daily intake in the same way for all foodstuffs. Contrary to this, cadmium absorption via tobacco smoking gives rise to very high blood cadmium levels which may not be accompanied by a proportionate increase in kidney burden (Elinder et al., 1983).

3.3 Health effects

Ingestion of highly contaminated food and drink may give rise to local gastrointestinal symptoms including vomiting, diarrhoea and, in severe cases, shock. Contamination of food may arise from cadmium-containing solders in water pipes, cooling or heating devices, or from dissolution of cadmium in pottery painted with cadmium-containing pigments. Formerly, the use of cadmium-plated cooking utensils was a common source of acute cadmium intoxication. Concentrations of 15 mg cadmium litre⁻¹ in water may give rise to acute symptoms including vomiting. Acute manifestations of cadmium intoxication arising from marine pollution have not occurred.

Chronic cadmium intoxication may be a result of long-term exposure via inhalation of cadmium fumes or dust, or from peroral exposure to contaminated food or beverages. The critical organ, i.e., the organ in which the first signs of adverse effects may be seen, is the kidney. The critical effect is a decrease in renal tubular reabsorption of proteins. One major sign of this effect is an increased urinary excretion of low molecular weight proteins, such as beta₂-microglobulin and retinol-binding proteins. A continuous catabolism of the cadmium metallothionein takes place after reabsorption, and cadmium is split from the metallothionein and bound to newly formed metallothionein in the tubular cells. It is supposed (Friberg et al., 1974; Nordberg, 1978; Nomiyama & Nomiyama, 1982) that kidney damage is prevented until such a stage is reached that the kidneys no longer produce enough metallothionein, and the free cadmium ions become very toxic to enzymatic processes. In more advanced cases, more extensive kidney damage occurs. This type of cadmium intoxication has frequently been observed following inhalation of cadmium in certain industries and after ingestion of contaminated food, particularly rice where, for example, contaminated water has been used for irrigation in certain areas of Japan. In contaminated areas, concentrations in certain crustaceans may also contribute significantly to high exposure levels. Once tubular proteinuria is manifest, it persists even though exposure ceases.

Other signs of cadmium intoxication which may occur at a later stage include anaemia and liver disorders. At a very late stage, effects on the bone in the form of osteoporosis and/or osteomalacia have been observed for cases of industrial exposure and in Japan as a result of ingestion of contaminated food. In Japan, the ensuing disease has been called Itai-itai, the manifestations of which are a combination of severe renal tubular damage and osteomalacia.

Even mild trauma may give rise to multiple fractures of the skeleton. The detailed pathogenesis of Itai-itai disease is not clear. Factors other than cadmium, such as low intake of calcium, proteins, and vitamin D, have been of importance, but cadmium is a necessary factor for the development of the disease. At present, the occurrence of

Some animal experiments indicate that hypertension can be induced by cadmium, but there are no convincing data indicating that cadmium can give rise to such symptoms in human beings.

Animal data show conclusively that injection of cadmium may cause sarcoma at the site of injection and also interstitial tumours of the testes. In a recent study, long-term inhalation of a cadmium chloride aerosol resulted in a pronounced and dose-related increase in the incidence of lung cancer (Takenaka et al., 1983). There is as yet no evidence from animal experiments which indicate that peroral exposure to cadmium increases the risk of developing cancer. Some epidemiological evidence exists, however, which suggests that cadmium may contribute to the development of cancer of the prostate as judged from studies on heavily-exposed workers (Belman & Nordberg, 1981). There are also some data suggesting a possible role of cadmium in the development of lung cancer in exposed workers. The recent animal data referred to above strengthen the suspicion that inhaled cadmium may be a human carcinogen for lung cancer. IARC (1976) concluded that occupational exposure to cadmium in some form, possibly the oxide, increases the risk of prostatic cancer and that one study also suggested an increase of respiratory cancer.

3.4 Total exposure to cadmium

Man is exposed to cadmium from the working environment, ambient air, drinking-water, tobacco, and food. For the non-occupationally exposed, food is the major source of intake. Among non-smokers, food could contribute 80 - 90% of the total intake of cadmium. In countries where intake via food is low, smoking is a major source of exposure and may contribute about half the body burden of cadmium. Normally, 0.1 - 0.2 µg cadmium is inhaled by smoking one cigarette. Concentrations of cadmium in ambient air are on an average approximately 5 ng m⁻³ and will not contribute significantly to the daily intake of cadmium.

Méranger et al. (1981) reported on cadmium concentrations in raw, treated, and distributed water from 71 municipalities in Canada. The mean cadmium concentration was ≤ 0.02 µg cadmium litre⁻¹ with a range of $\leq 0.02 - 0.9$ µg cadmium litre⁻¹.

In the absence of specific sources of contamination, most foodstuffs will contain less than 0.5 mg cadmium kg⁻¹ wet weight (Friberg et al., 1974; FAO/WHO, 1982; Elinder, in press). Low concentrations (1 - 50 µg kg⁻¹) are reported for meat, fish, and fruit. Somewhat higher concentrations have been reported for vegetables and cereal crops, e.g., wheat and rice, where concentrations in unpolluted areas may range from 0.01 to 0.15 mg cadmium kg⁻¹. In polluted areas, concentrations often reach 0.3 - 0.5 mg cadmium kg⁻¹. High concentrations (0.01 - 1 mg cadmium kg⁻¹) are found in the liver and kidneys of adult animals. Adult horses may contain up to 10 mg cadmium kg⁻¹ in liver and 10 - 150 mg cadmium kg⁻¹ in kidney cortex.

Data on total daily intake may also be unreliable due to a lack of quality assurance. Available data indicate, however, that the daily intake of cadmium in the USA and Europe is in the order of 20 µg cadmium, although large individual variations exist. In Japan, the average cadmium intake is 40 - 50 µg cadmium day⁻¹. In contaminated areas, the daily cadmium intake may be several times greater (Elinder, in press).

3.5 Contribution of cadmium from marine food

Data on cadmium concentrations in marine food indicate the general levels of exposure that may be expected. Most fish species contain relatively low cadmium levels of less than 0.4 µg g⁻¹. Some species, however, have somewhat higher concentrations; for example, flounder can contain up to 7 µg g⁻¹. For average per capita consumption of fish, however, this is a minor source of cadmium intake.

Certain species of mussels, scallops, and oysters often have cadmium concentrations exceeding 1 mg cadmium kg⁻¹ (FAO/WHO, 1982). In New Zealand oysters, cadmium concentrations ranging up to 8 mg cadmium kg⁻¹ wet weight have been found (Nielsen, 1975). Brown meat of crabs may contain 1 - 30 mg cadmium kg⁻¹ (UK Ministry of

determine the forms of cadmium in these organisms and their availability following ingestion by man. This source of cadmium could be of significance to total cadmium intake for particular regional population groups.

3.6 Evaluation of potential health effects

For most people, food is the major source of cadmium exposure. Special risk groups are smokers, who readily acquire blood cadmium levels which, on an average, are twice those in non-smokers. The average absorption of cadmium via food is probably around 5%, but with low body iron stores or calcium deficiency, absorption can increase up to 10 - 20%. There are, however, no large scale epidemiological studies to confirm the role of nutritional deficiencies. Analyses of cadmium in blood and kidney cortex in normal people show that 90 percentile values for kidney cortex levels are often about twice the median values. The difference between 90 percentile values and median values for blood levels is often greater. In some studies, the quotient is 5 or higher (Vahter, 1982; Friberg & Vahter, 1983).

Cadmium has been called the dissipated element because of its widespread occurrence in the environment and in different products. This implies the achievement of reduced cadmium exposure, measures have to be taken against a number of sources in order to minimize environmental contamination. For the average individual, uptake of cadmium in staple foods such as wheat and rice from contaminated soil is probably of greatest importance. For people with special food habits, other foods may be of equal or even greater importance. Thus, extensive long-term consumption of liver and kidney from certain animal species may considerably increase the daily intake of cadmium. Similarly, certain marine species, such as mussels and oysters, may contain high concentrations of cadmium and there are examples where extreme consumption of such oysters has led to a daily cadmium intake of 200 - 500 μg cadmium, which is more than 10 times the average daily intake of cadmium usually found.

In cadmium intoxication, the kidney cortex is the critical organ, the first signs being tubular dysfunction. The critical concentration is established on an individual level and varies among individuals. A WHO Task Group (WHO, 1977a), engaged in the preparation of WHO Environmental Health Criteria for Cadmium, has estimated the most likely critical concentration to be 200 $\mu\text{g g}^{-1}$ wet weight in kidney cortex. At that time, the variations among individuals were not taken into consideration. Friberg & Kjellström (1981) and Kjellström et al. (1984) proposed a "population critical concentration" (PCC) taking into consideration the distribution of the critical concentrations within a population. *In vivo* analyses of cadmium in the renal cortex of workers with and without cadmium-induced renal tubular damage have produced the following estimates of the average critical concentration: (PCC₅₀) of 332 $\mu\text{g cadmium g}^{-1}$ (Roels et al., 1983) and 319 $\mu\text{g cadmium g}^{-1}$ (Ellis et al., 1981). The individual variation is considerable in both studies, and Roels et al. (1938) estimated the PCC₁₀ at 216 $\mu\text{g cadmium g}^{-1}$ wet weight. It should be pointed out that the effects measured in these studies may not have been the earliest signs of cadmium intoxication. Furthermore, the empirical data referred to above by Ellis et al. (1981) and Roels et al. (1983) should be reduced by about 20%, since they used a factor of 1.5 in calculating concentrations in the kidney cortex from concentrations in the whole kidney. Recently, it has been observed that the factor should more likely be 1.25 (Svartengren et al., in press).

If 5% absorption of cadmium is assumed with 30% distributed to the kidneys, and a one-compartment metabolic model were used (half-time = 20 years), an intake of 200 - 400 $\mu\text{g cadmium day}^{-1}$ would result in approximately 200 $\mu\text{g cadmium g}^{-1}$ in the kidney cortex after approximately 50 years of exposure (Friberg et al., 1974). An absorption of 10% would correspondingly give 200 $\mu\text{g g}^{-1}$ in the kidney following an intake of 100 - 200 $\mu\text{g cadmium d}^{-1}$ over a 50-year period. Epidemiological data relating to exposure and effects after peroral intake are available from Japan. In the interim report by the above-mentioned WHO Task Group, it was estimated that, using the most sensitive analytical methods, a daily intake of approximately 200 $\mu\text{g cadmium day}^{-1}$ had caused the occurrence of an increased excretion of low molecular weight proteins. Thus, estimates based on critical concentrations in kidneys and metabolic models, together with epidemiological data, are in fairly close agreement.

4. Conclusions on Cadmium

4.1 Potential harm to living resources

Cadmium inputs to the marine environment are derived from both natural (volcanoes, dusts, and runoff) and anthropogenic sources. A major industrial use of cadmium is in electroplating, and discharges can occur from this activity. Other point source inputs are derived from zinc ore-smelting and phosphate fertilizer manufacture. Diffuse atmospheric inputs are derived from combustion of fossil fuels and domestic wastes. Estimates of oceanic fluxes of cadmium indicate that 50% is of anthropogenic origin, and this may have increased slightly in recent years. The normal range of cadmium in offshore water is 0.01 - 0.1 $\mu\text{g litre}^{-1}$; higher levels occur near point source inputs. Cadmium may be released into the water during dredging and disposal of dredged spoil, although the extent to which this occurs is not known.

Organocadmium compounds may be discharged to the marine environment. There is evidence, however, that alkyl compounds are rapidly degraded by abiotic processes in the environment.

Data from a large number of laboratory experiments show that a wide range of marine species are not acutely affected at cadmium concentrations less than 15 $\mu\text{g cadmium litre}^{-1}$. Moreover, very few of the species tested exhibited chronic effects from long-term exposure below this concentration, although one or two species of algae, two species of coelenterate, and one crustacean were harmed by concentrations within the range of 1 - 15 $\mu\text{g cadmium litre}^{-1}$; it is likely that there are other, untested species of similar sensitivity. However, it would appear that the present level of cadmium in offshore water is below that which would cause harmful effects to marine biota in general. Elevated concentrations in the vicinity of point source discharges could, however, cause harm to local species. Since estuaries and coastal waters are frequently the recipients of point source inputs, the human consumption of fish and shellfish from these areas may be affected either directly or indirectly.

Some resistant species, such as crabs and molluscs, may accumulate high levels of cadmium in their soft tissues without apparent detriment to their well-being. This accumulation may be derived from soluble cadmium in the water, or through the food chain via cadmium-contaminated algae, and this can occur even from low concentrations, in water not affected by discharges. The element can be complexed in tissues, but the availability of such cadmium to natural predators of these organisms is not known.

4.2 Potential hazards to human health

On an average, approximately 5% of ingested cadmium is absorbed, but individuals with a low iron store or on a calcium-deficient diet may have considerably higher absorption (up to 20%). About one third of the body burden of cadmium is located in the kidneys, where the concentration in kidney cortex is about 1.25 times the average concentration in the whole kidney. The biological half-time is long, approximately 20 years, and cadmium is, therefore, accumulated in the body. The total body burden at birth is only 1 μg , but continuous accumulation takes place in the body and, at age 50, the total body burden is between 10 and 30 mg, with concentrations in the kidney cortex of 15 - 50 $\mu\text{g cadmium g}^{-1}$ wet weight.

The first sign of cadmium intoxication is a dysfunction of the kidneys in the form of a decreased renal tubular absorption of proteins. This is the critical effect of cadmium with long-term exposures. If such an effect were prevented, even more serious effects could be avoided.

A WHO Task Group engaged in the preparation of the WHO Environmental Health Criteria for Cadmium has estimated the most likely critical concentration (when a small percentage of exposed people will show effects) to be 200 $\mu\text{g g}^{-1}$ wet weight in the kidney cortex. If a 5% absorption of cadmium is assumed, this concentration will be reached after an approximately 50-year exposure period to 200 - 400 $\mu\text{g cadmium day}^{-1}$. If a 10% absorption is assumed, the corresponding daily intake would be 100 - 200 $\mu\text{g cadmium day}^{-1}$.

order of 20 µg, although large variations among individuals exist. In Japan, the average cadmium intake is 40 - 50 µg day⁻¹. A Joint FAO/WHO Expert Committee in 1972 recommended a provisional tolerable weekly intake of 400 - 500 µg of cadmium (FAO/WHO, 1972).

Generally, low concentrations (less than 0.4 mg cadmium/kg) are reported for fish meat. Certain species of molluscs, scallops, and oysters, however, may often have cadmium concentrations exceeding 1 mg cadmium kg⁻¹. In New Zealand oysters, cadmium concentrations ranging up to 8 mg cadmium kg⁻¹ wet weight have been found. The brown meat of crabs may contain 1 - 30 mg cadmium kg⁻¹ wet weight.

Assuming that an average fish eater consumes 40 g of fish per day (actual fish consumption values are given in Table 1), cadmium in fish will constitute only a small fraction of the total daily intake. Only under the most exceptional circumstances (very high consumption and very high cadmium levels in fish) will the cadmium intake from fish constitute an important part of the total daily intake via food. It is possible to find examples, however, where the high consumption of certain shellfish may considerably increase the total daily intake of cadmium. It has been reported recently that people consuming large numbers of certain New Zealand oysters have a weekly intake of 1400 - 3500 µg cadmium, which is considerably above the provisional tolerable weekly intake.

5. References

- ALABASTER, J.S. (1978) Ecotoxicity of cadmium, Stevenage, United Kingdom, Water Research Centre, 38 pp.
- ASTON, S.R., CHESTER, R., & GRIFFITHS, A. (1972) Distribution of cadmium in north Atlantic deep-sea sediments. Nature (London), 239: 393.
- AYLETT, B.J. (1973) The chemistry of Zn, Cd, Hg, Oxford, Pergamon Press, Vol. 18 (Pergamon Texts in Inorganic Chemistry).
- BELMAN, S. & NORDBERG, G.F., ed. (1981) Workshop/Conference on the role of metals in carcinogenesis, Atlanta, Georgia, March 24-28, 1980. Environ. Health Perspect., 40: 1-42.
- BEWERS, J.M. & YEATS, P.C. (1981) Behaviour of trace metals during estuarine mixing. In: River inputs to ocean systems. UNEP/IOC/SCOR Review Workshop, Rome, March, 1979, New York, United Nations, pp. 103-115.
- BUAT-MENARD, P. & ARNOLD, M. (1978) The heavy metal chemistry of atmospheric particulate matter emitted by Mount Etna volcano. Geophys. Res. Lett., 5: 245-248.
- BULL, K.R., MURTON, R.K., OSBORN, D., WARD, P., & CHENG, L. (1977) High levels of Cd in Atlantic seabirds and sea-skaters. Nature (London), 269: 507-509.
- CALABRESE, A., THURBERG, F.P., DAWSON, M.A., & WENZLOF, D.R. (1975) Sub-lethal physiological stress induced by cadmium and mercury in the winter flounder Pseudopleuronectes americanus. In: Koeman, J.H. & Strik, J.J.T.W.A., ed. Sublethal effects of toxic chemicals on aquatic animals, Amsterdam, Elsevier, pp.15-21.
- CAMNER, P., CLARKSON, T.W., & NORDBERG, G.F. (1979) Routes of exposure, dose and metabolism of metals. In: Friberg, L., Nordberg, G.F., & Vouk, V.B., ed. Handbook on the toxicology of metals, Amsterdam, Elsevier, pp. 65-97.
- CAMPBELL, J.A. & LORING, D.H. (1980) Baseline levels of heavy metals in waters and sediments of Baffin Bay. Mar. Pollut. Bull., 11: 257-261.
- CARMICHAEL, N.G. & FOWLER, B.A. (1981) Cadmium accumulation and toxicity in the kidney of the Bay scallop Argopectan irradians. Mar. Biol., 65: 35-43.
- CARPENE, E. & GEORGE, S.G. (1981) Absorption of cadmium by gills of Mytilus edulis L. Mol. Physiol., 1: 23-34.
- CHERIAN, M.G. & SHATKH, Z.A. (1975) Metabolism of intravenously injected

CHERIAN, M.G., GOYER, R.A., & VALBERG, L.S. (1978) Gastrointestinal absorption and organ distribution of oral cadmium chloride and cadmium-metallothionein in mice. J. Toxicol. environ. Health, 4: 861-868.

COLEMAN, N. (1980) The effect of emersion on cadmium accumulation by Mytilus edulis. Mar. Pollut. Bull., 11: 359-362.

COOMBS, T.L. (1979) Cadmium in aquatic organisms. In: Webb, M., ed. The chemistry, biochemistry, and biology of cadmium, Amsterdam, Elsevier, pp. 93-139.

COOMBS, T.L. (1983) Personal communication, Aberdeen, Scotland, NERC Institute of Marine Biochemistry.

DAVIES, I.M. (1981) Survey of trace elements in fish and shellfish landed at Scottish Ports, 1975-76, 28 pp (DAFS Scottish Fisheries Research Report No. 19) (Department of Agriculture and Fisheries for Scotland).

DAWSON, M.A., GOULD, E., THURBERG, F.P., & CALABRESE, A. (1977) Physiological responses of juvenile striped bass, Morone saxatilis to low levels cadmium and mercury. Chesapeake Sci., 8: 353-359.

DENTON, G.R.W. & BURDEN-JONES, C. (1981) Influence of temperature and salinity on the uptake, distribution, and depuration of mercury, cadmium, and lead by the Black-lip oyster Saccostrea echinata. Mar. Biol., 64: 317-326.

DUCE, R.A., HOFFMAN, G.L., AND OTHERS (1976) Trace metals in the marine atmosphere, sources, and fluxes. In: Windom, H.L. & Duce, R.A., ed. Marine pollutant transfer, Massachusetts, Lexington Books, pp. 77-119.

ELINDER, C.-G. (in press) Cadmium: uses, occurrence, and intake. In: Friberg, L., Elinder, C.-G., Kjellström, T., & Nordberg, G.F., ed. Cadmium and health, Boca Raton, Florida, CRC Press.

ELINDER, C.-G., KJELLSTROM, T., FRIBERG, L., LIND, B., & LINNMAN, L. (1976) Cadmium in kidney cortex, liver, and pancreas among Swedish autopsies. Arch. environ. Health, 31: 292-302.

ELINDER, C.-G., FRIBERG, L., LIND, B., & JAWAID, M. (1983) Lead and cadmium levels in blood samples from the general population of Sweden. Environ. Res., 30: 233-253.

ELLIS, K.J., MORGAN, W.D., ZANZI, I., YASUMURA, S., VARTSKY, D., & COHN, S.H. (1981) Critical concentrations of cadmium in human renal cortex: dose-effect studies in cadmium smelter workers. J. Toxicol. environ. Health, 7: 691-703.

FALCONER, C.R., DAVIES, I.M., & TOPPING, G. (1983) Trace metals in the common porpoise, Phocoena phocoena. Mar. environ. Res., 8: 119-127.

FAO/WHO (1972) Evaluation of certain food additives and the contaminants mercury, lead, and cadmium. Sixteenth report of the Joint FAO/WHO Expert Committee on Food Additives, Geneva, World Health Organization (Technical Report Series No. 505).

FAO/WHO (1982) GEMS: Global Environmental Monitoring System, Joint FAO/WHO Food and Animal Feed Contamination Monitoring Programme, Summary of Data received from Collaborating Centres for Food Contamination Monitoring, Geneva, World Health Organization.

FASSETT, D.W. (1980) Cadmium. In: Waldron, H.A., ed. Metals in the environment, London, Academic Press, pp. 61-110.

FISHER, N.S. & FROUD, D. (1980) Heavy metals and marine diatoms. Influence of dissolved organic compounds in toxicity and selection for metal tolerance among four species. Mar. Biol., 59: 85-93.

FLEISCHER, M., SAROFIM, A.F., FASSETT, D.W., HAMMOND, P., SHACKLETTE, H.T., NISBET, I.C.T., & EPSTEIN, S. (1974) Environmental impact of cadmium: a review by the panel on hazardous trace substances. Environ. Health Perspect., 7: 253-323.

FORSTNER, U. & VAN LIERDE, J.H. (1979) Trace metals in water purification processes. In: Förstner, U. & Wittman, G., ed. Metal pollution the marine environment, Berlin, Springer-Verlag, Chapter G, pp. 324-359.

FREI, R.W. & HUTZINGER, O., ed. (1976) Analytical aspects of mercury and other metals in the environment, New York, Gordon and Breach, 210 pp.

FRIBERG, L.T. & KJELLSTROM, T. (1981) Toxic metals: pitfalls in risk estimation. In: Proceedings of the International Conference on Heavy Metals in the Environment, Amsterdam, September, 1981, Edinburgh, CEP Consultants Ltd, pp. 1-11.

FRIBERG, L.T. & VAHTER, M. (1983) Assessment of exposure to lead and cadmium through biological monitoring. Results of a UNEP/WHO global study. Environ. Res., 30: 95-128.

FRIBERG, L.T., PISCATOR, M., NORDBERG, G.F., & KJELLSTROM, T. (1974) Cadmium in the environment, Cleveland, Ohio, CRC Press.

FRIBERG, L.T., ELINDER, C.-G., KJELLSTROM, T., & NORDBERG, G.F., ed. (in press) Cadmium and health, Boca Raton, Florida, CRC Press.

GENTILE, S.M., GENTILE, J.H., WALKER, J., & HELTSHE, J.F. (1982) Chronic effects of cadmium on two species of mysid shrimp: Mysidopsis bahia and Mysidopsis bigelowi. Hydrobiologia, 93: 195-204.

GEORGE, S.G. (1980) Correlation of metal accumulation in mussels with the mechanisms of uptake, metabolism, and detoxification: a review. Thalassia Jugosl., 16: 347- 365.

GEORGE, S.G. & COOMBS, T.L. (1977) The effects of chelating agents on the uptake and accumulation of cadmium by Mytilus edulis. Mar. Biol., 39: 261-268.

GEORGE, S.G. & PIRIE, B.J.S. (1979) The occurrence of cadmium in subcellular particles in the kidney of the marine mussel, Mytilus edulis exposed to cadmium. Biochim. Biophys. Acta, 580: 234-244.

GOULD, E. (1977) Alteration of enzymes in winter flounder, Pseudopleuronectes americanus, exposed to sub-lethal amounts of cadmium chloride. In: Vernberg, F.J., Calabrese, A., Thurberg, F.P., & Vernberg, W.B., ed. Physiological responses of marine biota to pollutants, New York, Academic Press, pp. 209-224.

GOULD, E. (1980) Low salinity stress in the American lobster, Homarus americanus after chronic sublethal exposure to cadmium: biochemical effects. Helgolander Meeresunters., 33: 36-46.

HARDY, J.T., SCHMIDT, R.L., & APTS, C.W. (1981) Marine sediment and interstitial water: effects on bioavailability of cadmium of the gills of the clam Protothaca staminea. Bull. environ. Contam. Toxicol., 27: 798-805.

HUTTON, M. (1982) Cadmium in the european community, London, Chelsea College, Monitoring and Assessment Research Centre, 99 pp (MARC Report No. 26).

IARC (1976) The evaluation of carcinogenic risk of chemicals to humans, Lyons, International Agency for Research on Cancer (IARC Monographs Vol. 11).

IRPTC (1981) Data profile on cadmium in the marine and estuarine environment, Geneva, International Register for Potentially Toxic Chemicals (UNEP).

JANSSEN, H.H. & SCHOLZ, N. (1979) Uptake and cellular distribution of cadmium in Mytilus edulis. Mar. Biol., 55: 133-141.

KAYSER, H. (1982) Cadmium effects in food chain experiments with marine plankton algae (Dinophyta) and benthic filter feeders (Tunicata). Netherlands J. Sea Res., 16: 444-454.

- KAYSER, H. & SPERLING, K.R. (1980) Cadmium effects and accumulation in cultures of Prorocentrum micans (Dinophyta). Helgolander Meeresunters., 33: 89-102.
- KJELLSTROM, T. (1979) Exposure and accumulation of cadmium in populations from Japan, the United States, and Sweden. Environ. Health Perspect., 28: 169-197.
- KJELLSTROM, T. & NORDBERG, G.F. (1978) A kinetic model of cadmium metabolism in the human being. Environ. Res., 16: 248-269.
- KJELLSTROM, T., ELINDER, C.-G., & FRIBERG, L.T. (1984) Conceptual problems in establishing the critical concentration of cadmium in human kidney cortex. Environ. Res., 33: 284-295.
- KOHLER, K. & RIISGARD, H.U. (1982) Formation of metallothioneins in relation to accumulation of cadmium in the common mussel Mytilus edulis. Mar. Biol., 66: 53-59.
- KREMLING, K., PIUZE, J., VON BROCKEL, K., & WONG, C.S. (1978) Studies on the pathways and effects of cadmium in marine plankton communities in experimental enclosures. Mar. Biol., 48: 1-10.
- KRENKEL, P.A., HARRISON, J., & BURDICK, J.C., ed. (1976) Dredging and its environmental effects, New York, American Society of Civil Engineers, 1035 pp.
- KUIPER, J. (1981) Fate and effects of cadmium in marine plankton communities in experimental enclosures. Mar. Ecol. Prog. Ser., 6: 161-174.
- LARSSON, A. (1975) Some biochemical effects of cadmium in fish. In: Koeman, J.H. & Strick, J.J.T.W.A., ed. Sublethal effects of toxic chemicals on aquatic animals, Amsterdam, Elsevier, pp. 3-13.
- LARSSON, A., BENGTSSON, B.-E., & HAUX, C. (1981) Disturbed ion balance in flounder, Platichthys flesus L. exposed to sublethal levels of cadmium. Aquat. Toxicol., 1: 19-35.
- LI, W.K.W. (1980) Cellular accumulation and distribution of cadmium in Isochrysis galbana during growth inhibition and recovery. J. Plankton Res., 2: 283-294.
- MACKAY (1983) Metal organic complexes in sea water. An investigation of naturally-occurring complexes of Cu, Zn, Fe, Mn, Ni, Mg, and Cd using high performance liquid chromatography with atomic fluorescence detection. Mar. Chem., 13: 169-180.
- MAGNUSSON, B. & WESTERLUND, S. (1980) The determination of Cd, Cu, Fe, Ni, Pb, and Zn in Baltic Sea water. Mar. Chem., 8: 231-244.
- MARKHAM, K.W., KREMER, B.P., & SPERLING, K.R. (1980) Effects of cadmium on Laminaria saccharina in culture. Mar. Ecol. Res. Ser., 3: 31-39.
- MARSHALL, A.T. & TALBOT, V. (1979) Accumulation of cadmium and lead in the gills of Mytilus edulis: X-ray microanalysis and chemical analysis. Chem-Biol. Interact., 27: 111-123.
- MCKENZIE, J., KJELLSTROM, T., & SHARMA, R.P. (1982) Cadmium intake, metabolism, and effects in people with a high intake of oysters in New Zealand, Washington (Report to US Environmental Protection Agency (Grant ID No. R807058-01-0)).
- MCKIE, J.C., DAVIES, I.M., & TOPPING, G. (1980) Heavy metals in Grey Seals (Halichoerus grypus) from the east coast of Scotland. International Council for the Exploration of the Sea, 13 pp (CM 1980, E.41).
- MCLEESE, D.W. (1980) Uptake and excretion of cadmium by marine organisms from sea water with cadmium at low concentration: a review. In: Uthe, J.F. & Zitko, V., ed. Cadmium pollution of Belledune Harbour, New Brunswick, Canada. Can. Tech. Rep. Fish Aquat. Sci., 963: 55-63.
- MERANGER, J.C., SUBRAMANIAN, K.S., & CHALIFOUX, C. (1981) Metals and other elements:

- MIRKES, D.Z., VERNBERG, W.B., & DE COURSEY, P.J. (1978) Effects of cadmium and mercury on the behavioural responses and development of Eurypassopeus depressus larvae. Mar. Biol., 47: 143-147.
- MOHLENBERG, F. & JENSEN, A. (1980) The ecotoxicology of Cd in fresh and sea water and water pollution with Cd in Denmark, Charlottenlund, Denmark, 42 pp, National Agency of Environmental Protection.
- MOWDY, D.E. (1981) Elimination of laboratory-acquired cadmium by the oyster, Crassostrea virginica in the natural environment. Bull. environ. Contam. Toxicol., 26: 345-351.
- MURRAY, A.J. (1979) Metals, organochlorine pesticide, and PCB residue in fish and shellfish landed in England and Wales during 1974. Aquat. Environ. Monit. Rep., 2: 11.
- MURRAY, L.A. & NORTON, M.G. (1979) The composition of dredged spoils dumped at sea from England and Wales. MAFF Fish. Res. Tech. Rep., 52: 10.
- MURRAY, L.A., NORTON, M.G., NUNNY, R.S., & ROLFE, M.S. (1980) The field assessment of effects of dumping wastes at sea. VI. The disposal of sewage sludge and industrial waste off the River Humber. MAFF Fish. Res. Tech. Rep., 55: 35.
- NICHOLSON, R.A. & MOORE, P.J. (1981) The distribution of heavy metals in the superficial sediments of the North Sea. Rapp. P.-v. Reun. Cons. int. Explor. Mer., 181: 35-48.
- NIELSEN, S.A. (1975) Cadmium in New Zealand dredge oysters: geographic distribution. Int. J. environ. anal. Chem., 4: 1-7.
- NIMMO, D.R., RIGBY, R.A., BAHNER, L.H., & SHEPPARD, J.M. (1978) The acute and chronic effects of cadmium on the estuarine mysid, Mysidopsis bahia. Bull. environ. Contam. Toxicol., 19: 80-85.
- NOEL-LAMBOT, F. (1981) Presence in the intestinal lumen of marine fish of corpuscles with a high cadmium-, zinc-, and copper-binding capacity: a possible mechanism of heavy metal tolerance. Mar. Ecol. Prog. Ser., 4: 175-181.
- NOMIYAMA, K. & NOMIYAMA, H. (1982) Tissue metallothioneins in rabbits chronically exposed to cadmium, with special reference to the critical concentration of cadmium in the renal cortex. In: Foulkes, E.C., ed. Biological roles of metallothionein, New York, Elsevier, pp. 47-67.
- NORDBERG, G.F., GOYER, R.A., & NORDBERG, M. (1975) Comparative toxicity of cadmium-metallothionein and cadmium chloride on mouse kidney. Arch. Pathol., 99: 192-197.
- NORDBERG, M. (1978) Studies on metallothionein and cadmium. Environ. Res., 15: 381-404.
- NRIAGU, J.O., ed. (1980) Cadmium in the environment. Part. I: Ecological cycling, Toronto, Wiley-Interscience.
- OJAVEER, E., ANNIST, J., JANKOWSKI, H., PALM, T., & RAID, T. (1980) On effect of copper, cadmium, and zinc on the embryonic development of Baltic spring spawning herring. Finn. Mar. Res., 247: 135-140.
- PAGANO, G., ESPOSITO, A., & GIORDANO, G.G. (1982) Fertilization and larval development in sea urchins following exposure of gametes and embryos to cadmium. Arch. environ. Contam. Toxicol., 11: 47-55.
- PAGE, A.L., BINGHAM, F.T., & CHANG, A.C. (1981) Cadmium. In: Lepp, N.W., ed. Effect of heavy metal pollution on plants, London, Applied Science, Vol. 1, pp. 77-109 (Pollution Monitoring Series).
- PARIS COMMISSION (1983) Third Annual Report of the Paris Commission, London.

- PETERSON, P.J. & ALLOWAY, B.J. (1979) Cadmium in soils and vegetation. In: Webb, M., ed. The chemistry, biochemistry, and biology of cadmium, Amsterdam, Elsevier, pp. 45-92.
- POULSON, E., RIISGARD, H.U., & MOHLENBERG, F. (1982) Accumulation of cadmium and bioenergetics in the mussel Mytilus edulis. Mar. Biol., 68: 25-29.
- PRESTON, A., JEFFERIES, D.F., DUTTON, J.W.R., HARVEY, B.R., & STEELE, A.K. (1972) British Isles coastal waters: the concentrations of selected heavy metals in sea water, suspended matter, and biological indicators: a pilot survey. Environ. Pollut., 3: 69-82.
- PREVOT, P. (1980) Etude de l'action intracellulaire de cadmium. Observation et mesures d'absorption. In: Les métaux en milieu marin, Paris, Vol. 2, pp. 117-133 (Editions du Centre Nationale de la Recherche Scientifique).
- PRICE, R.K.J. & UGLOW, R.F. (1980) Cardiac and ventilatory responses of Crangon crangon to cadmium, copper, and zinc. Helgolander Meeresunters., 33: 59-67.
- RAHOLA, T., AARAN, R.-K., & MIETTINEN, J.K. (1972) Half-time studies of mercury and cadmium by whole-body counting. In: Assessment of Radioactive Contamination in Man. International Atomic Energy Agency Proceeding Series, New York, Unipublishers, pp. 553-562 (IAEA-SM 150/13).
- RAY, S., MCLEESE, D.W., & PEZZACK, D. (1980) Accumulation of cadmium by Nereis virens. Arch. environ. Contam. Toxicol., 9: 1-8.
- RAY, S., MCLEESE, D.W., & BURRIDGE, L.E. (1981a) Cadmium in tissues of lobsters captured near a lead smelter. Mar. Pollut. Bull., 12: 383-386.
- RAY, S., MCLEESE, D.W., & PETERSON, M.R. (1981b) Accumulation of copper, zinc, cadmium, and lead from two contaminated sediments by three marine invertebrates: a laboratory study. Bull. environ. Contam. Toxicol., 26: 315-322.
- ROELS, H., LAUWERYS, R., & DARDENNE, A.N. (1983) The critical level of cadmium in human renal cortex: a re-evaluation. Toxicol. Lett., 15: 357-360.
- ROJAHN, T. (1972) Determination of copper, lead, cadmium, and zinc in estuarine water by anodic-stripping alternating-current voltammetry on the hanging mercury drop electrode. Anal. Chim. Acta, 62: 438-441.
- RUBINSTEIN, N.I., LORES, E., & GREGORY, N.R. (1983) Accumulation of PCBs, mercury, and cadmium by Nereis virens, Mercenaria mercenaria, and Palaemonetes pugio from contaminated harbour sediments. Aquat. Toxicol. (in press).
- SCHULZ-BALDES, M. & CHENG, L. (1980) Cadmium in Halobates micans from the central and south Atlantic Ocean. Mar. Biol. (Berlin), 59: 163-168.
- SIMPSON, W.R. (1981) A critical review of cadmium in the marine environment. Prog. Oceanogr., 10: 1-70.
- SOMMERS, L.E. (1977) Chemical composition of sewage sludges and analysis of their potential use as fertilizers. J. environ. Qual., 6: 225-232.
- SOMMERS, L.E., TABATABAI, M.A., & NELSON, D.W. (1977) Forms of sulphur in sewage sludge. J. environ. Qual., 6: 42-46.
- SOYER, M. & PREVOT, P. (1981) Ultrastructural damage by cadmium in a marine dinoflagellate, Prorocentrum micans. J. Protozool., 28: 308-313.
- STOVER, R.C., SOMMERS, L.E., & SILVIERA, D.J. (1976) Evaluation of metals in waste water sludge. J. Water Pollut. Res. Fed., 48: 2165-2175.
- STROMGREN, T. (1982) Effects of heavy metals (Zn, Hg, Cu, Cd, Pb, Ni) on the length growth of Mytilus edulis. Mar. Biol. (Berlin), 72: 69-72.

TAKENAKA, S., OLDIGES, H., KONIG, H., HOCHRAINER, D., & OBERDORSTER, G. (1983) Carcinogenicity of cadmium chloride aerosols in Wistar rats. J. Natl Cancer Res. Inst., 70: 367-373.

TAYLOR, D. (1981) A summary of the data on the toxicity of various materials to aquatic life. II. Cadmium, 2nd ed., Brixham, Devon, United Kingdom, ICI Ltd, 50 pp (Report No. BL/A/2073).

THEEDE, H., SCHOLZ, N., & FISCHER, H. (1979) Temperature and salinity effects on the acute toxicity of cadmium to Laomedae loveni (Hydrozoa). Mar. Ecol. Prog. Ser., 1: 13-19.

THURBERG, F.P., CALABRESE, A., GOULD, E., GREIG, R.A., DAWSON, M.A., & TUCKER, R.K. (1977) Response of the lobster, Homarus americanus to sublethal levels of cadmium and mercury. In: Vernberg, F.J., Calabrese, A., Thurberg, F.P., & Vernberg, W.B., ed. Physiological responses of marine biota to pollutants, New York, Academic Press, pp. 185-207.

TSUCHIYA, K., ed. (1978) Cadmium studies in Japan: a review, Amsterdam, Elsevier.

UK MINISTRY OF AGRICULTURE, FISHERIES AND FOOD (1973) Survey of cadmium food. Working Party on the Monitoring of Foodstuffs for Heavy Metals, fourth report, London, Her Majesty's Stationery Office.

US EPA (1980) Ambient water quality criteria for cadmium, Washington DC, US Environmental Protection Agency, Criteria and Standards Division, 192 pp (US EPA No. 440/5-80-025).

VAHTER, M., ed. (1982) Assessment of human exposure to lead and cadmium through biological monitoring, Stockholm, Sweden, National Swedish Institute of Environmental Medicine and Karolinska Institute, Department of the Environmental Hygiene (Prepared for United Nations Environment Programme and WHO).

VON BIAS, R. (1981) Cadmium uptake rates in euryhaline amphipods of the Elbe-Estuary: experiments with Corophium volutator (Pallas) (Amphipoda, Corophiidae). Arch. Hydrobiol., 61: 84-152.

VON WESTERNHAGEN, H., DETHLEFSEN, V., ROSENTHAL, H., FURSTENBERG, G., & KLINCKMAN, J. (1978) Fate and effects of cadmium in a marine ecosystem. Helgolander wiss. Meeresunters., 31: 471-484.

VON WESTERNHAGEN, H., DETHLEFSEN, V., & ROSENTHAL, H. (1980) Correlation between cadmium concentration in the water and tissue residue levels in dab, Limanda limanda L., and plaice, Pleuronectes platessa L. J. Mar. Biol. Assoc. UK, 60: 45-58.

WARD, T.J. (1982) Effect of cadmium on particle clearance by the Sydney rock oyster, Saccostrea commercialis. Aust. J. Mar. Freshw. Res., 33: 711-715.

WATLING, H.R. (1982) Comparative study on the effects of zinc, cadmium, and copper on the larval growth of three oyster species. Bull. environ. Contam. Toxicol., 28: 195-201.

WEBB, M., ed. (1979) The chemistry, biochemistry, and biology of cadmium, Amsterdam, Elsevier.

WEIS, P. & WEIS, J.S. (1976) Effects of heavy metals on fin regeneration in the killifish, Fundulus heteroclitus. Bull. environ. Contam. Toxicol., 16: 197-202.

WHITE, S.L. & RAINBOW, P.S. (1982) Regulation and accumulation of copper, zinc, and cadmium by the shrimp Palaemon elegans. Mar. Ecol. Prog. Ser., 8: 95-101.

WHITFIELD, M., TURNER, D.A., & DICKSON, A. G. (1981) Speciation of dissolved constituents in estuaries. In: River Inputs to Ocean Systems, Proceedings Review Workshop, FAO, Rome, 26-30 March 1979, New York, UNEP/UNESCO, pp. 132-151.

WHO (1977) WHO Environmental Health Criteria for Cadmium. Ambio 6: 287-290

WHO (1979) Environmental Health Criteria for Cadmium, Geneva, World Health Organization, 111 pp (Interim Report No. EHE/EHC/79.20).

WHO (1980) Recommended health-based limits in occupational exposure to heavy metals, Geneva, World Health Organization (Report of a WHO study group) (Technical Report Series No. 647).

WITTMANN, G. (1979) Toxic metals. In: Förstner, U. & Wittmann, G., ed. Metal pollution in the aquatic environment, Berlin, Springer-Verlag, Chapter B, pp. 3- 70.

YEN-WAN, HUNG (1982) Effects of temperature and chelating agents on cadmium uptake in the American oyster. Bull. environ. Contam. Toxicol., 28: 546-551.

ZAROOGIAN, G.E. (1980) Crassostrea virginica as an indicator of cadmium pollution. Mar. Biol., 58: 275-284.

ZAROOGIAN, G.E. & MORRISON, G. (1981) Effect on cadmium body burdens in adult Crassostrea virginica on fecundity and viability of larvae. Bull. environ. Contam. Toxicol., 27: 344-348.

III. LEAD

1. Lead in the Marine Environment

1.1 Reference documentation

The major reviews consulted in the preparation of this section were The Biogeochemistry of Lead in the Environment (Nriagu, 1978), Lead in the Marine Environment (Branica & Konrad, 1980), and Lead: Environmental Health Criteria (WHO, 1977). Information on sources of lead contamination in the terrestrial environment was obtained from Friedman & Hutchinson (1981). Other papers that were used are listed in the reference section.

1.2 General facts

Lead has the chemical symbol Pb (Latin: plumbum). It is possibly the first metal discovered and worked, and it is the heaviest element in the Periodic Table Group IVb. Its atomic number is 82, and the atomic weight is 207.19 (depending on source). Lead has 2 oxidation states, Pb^{2+} and Pb^{4+} , and Pb^{2+} greatly predominates in the aquatic environment. In clean-fresh water at pH 9, $PbCO_3$ is the main inorganic species (88%); the remainder is $Pb(OH)_2$. At pH 6, $PbCO_3$ (15%), $PbSO_4$ (3%), and $PbCl_2$ (1%) are the prevailing ligands. In sea water, $PbCl_2$ (43%), $PbCO_3$ (42%), and $Pb(OH)_2$ (9%) (Whitfield et al., 1981) are found, but different authorities give varying speciation and ratios.

In aqueous solution, Pb^{2+} is a borderline Type (a) cation. Under appropriate conditions, alkyl-lead compounds can be formed in the environment (Harrison & Laxen, 1978), and this has been demonstrated in the laboratory (Wong, 1975; Chau & Wong, 1980), but these organo-metals may not be stable (Wood, 1980). Various lead sulphides are formed under anaerobic conditions in sediments.

"High lead" can occur as a result of one or more factors. These include Pb-mineralization, low pH from rock characteristics or the presence of organic acids arising from peat or tree cover, chelation, high chloride, bicarbonate, or nitrate content, intrusion of thermally active water, and/or deficiency of alkaline minerals. "Low lead" can result from Pb-deficient mineralization, a pH > 7, pressure of carbonate, agricultural application of lime, temporary uptake by profuse aquatic vegetation, and/or contact with marl, chalk, or other soils or sediments rich in alkaline minerals. Efficient removal of Pb from raw water requires extra processes carefully matched to the water characteristics, since routine purification processes are often ineffective.

1.3 Sources

Rocks containing small amounts of lead are common and widespread. Typical concentrations range from 10 to 20 $\mu g g^{-1}$ in many igneous and metamorphic rocks, from 10 to 70 $\mu g g^{-1}$ in carbonaceous shales, and about 100 $\mu g g^{-1}$ or more in some phosphate rocks.

Lead is recovered commercially from a range of locally-occurring ores of which galena (PbS) and, to a lesser extent, cerrusite ($PbCO_3$), anglesite ($PbSO_4$), and others are important. Often, deposits are also rich in zinc and zinc-copper. These mixed ores give us significant amounts of silver, gold, bismuth, antimony, arsenic, cadmium, tin, gallium, indium, germanium, and tellurium. Sources of lead are found in many countries of the world, and some ores (especially galena) are of high purity. The most important lead mining sites (i.e., those producing more than 10^5 tonnes year⁻¹) are found in Australia (10% of global production), Bulgaria (3%), Canada (9.6%), China (3.8%), Mexico (4.5%), Peru (5.5%), the USA (16%), the USSR (14.5%), and Yugoslavia (3.5%). In addition, 15% of the mining production is distributed on a minor scale among about 50 other countries. Substantial amounts of lead are recycled (estimated to be about 33% for some countries) and reclaimed from waste recovery processes. Global lead production in 1975 was 3.6×10^6 tonnes (all these data from WHO, 1977b and Koepe, 1977).

processes can give rise to water and air pollution, and further pollution is derived from the manufacture, use, and discarding of goods, alloys, and organo-lead compounds. Lead is released into the environment by the combustion of coal, wood, and other organic matter including city garbage (DOE, 1974).

Synthetic alkyl-lead fuel additives are produced on a large but slowly declining scale, (1973: 378 000 tonnes; 1974: 357 000 tonnes; 1975: 301 000 tonnes). Those additives are widely regarded as a serious source of atmospheric lead pollution. Efforts are being made to progressively reduce and, when possible, eliminate world-wide use of lead as a motor fuel additive. In 1969, the proportion of alkyl-lead to total lead consumed was 20% in the USA, 11% in the United Kingdom, 11% in Italy, and 5% in France, other countries remaining within this range of consumption.

Of much current concern is the use of lead in drinking-water distribution and domestic plumbing. This lead usage contributes to water pollution in rivers, estuaries, and the sea. Lead is a common contaminant of sewage wastes. As far as is known, lead, a potentially toxic and highly available element, is generally not essential to man or living organisms.

Nriagu (1978) estimates that the global atmospheric lead emissions from man-made sources were 438×10^3 tonnes year⁻¹ for the years 1974-75, in contrast to only 18.55×10^3 tonnes year⁻¹ from natural mobilization and inputs. Atmospheric discharges from the combustion of lead in petrol (61%), from steel and base metal production (23%), and from the mining and smelting of lead (8%) are the major contributors, with coal combustion and numerous minor activities comprising the remainder.

The main industrial uses of lead are summarized in Table 6. Many of these promote widespread dispersal of lead as do, for example, the emissions from leaded petrol and coal combustion; to this must be added other uses such as lead solders in canning. The recycling of metallic lead from domestic and industrial uses, such as in lead-acid batteries, removes a significant environmental threat.

Table 6. Lead consumption by use in Europe, the USA, and Japan
(x 1000 metric tonnes)^a

Use	1964	1969	1974	1980 (estimated)
Batteries	772	992	1390	1700
Pigments, chemicals	248	289	360	500
Tetraethylleads	254	319	317	200
Alloys	303	307	288	300
Cable sheathing	427	352	322	200
Pipe and sheet	298	266	173	200
Miscellaneous	206	192	258	200
Total	2508	2717	3108	3300

^a Sources: International Lead-Zinc Study Group, World Bureau of Metal Statistics; Lead Industries Association; US Bureau of Mines; Noranda Sales Corporation. From: Robinson (1978).

In addition to the many uses of lead metal and alloys, there are innumerable

hazard involved, lead pigments are no longer used in interior paints, but remain a component of some exterior paints and protective coatings for metals. Alkyl-lead antiknock derivatives are produced on a massive scale and are a cause of environmental concern; other organoleads used in biocides, detonators, plastics, and catalysts are produced on a minor scale and, biocides excepted, have less access to the environment.

There is a wide range of lead concentrations in the soil (2 - 200 $\mu\text{g g}^{-1}$) with considerable areal heterogeneity. Some geologically-unusual soils from diverse countries contain up to 30000 $\mu\text{g g}^{-1}$ (Nriagu, 1978). Usually, lead in the soil is virtually immobile and barely soluble, and lead in drainage waters is readily adsorbed by hydrous metal oxides, clay minerals, and organic-suspended particles.

Lead in rivers is found in the form of poorly soluble species together with many complexes such as organic acids, amino acids (often complexed with cysteine, which is exceptionally stable), and colloidal ones combined with peptides, proteins, and other natural macromolecules. Treated or untreated sewage residues and sewage sludge all possess enhanced levels of diverse organic lead compounds. Soluble lead concentrations vary widely, but typical ranges for the world's major rivers are 1 - 10 $\mu\text{g litre}^{-1}$, 1 - 55 $\mu\text{g litre}^{-1}$ for upland streams, 0.5 - 180 $\mu\text{g litre}^{-1}$ for rivers and streams in populated areas, with mineralized acid streams having upwards of 1000 $\mu\text{g litre}^{-1}$, and for run-off water from city centres 100 - 12 000 $\mu\text{g litre}^{-1}$, of which about 10% may be alkyl leads (Chow, 1978).

The lead concentration in river sediments is < 10 $\mu\text{g g}^{-1}$ for remote Arctic regions, an average of 23 $\mu\text{g g}^{-1}$ for those of the world's major rivers and a range of 50 - 500 $\mu\text{g g}^{-1}$ for large rivers traversing populous areas which rises to 3800 $\mu\text{g g}^{-1}$ for heavily-mineralized streams (Nriagu, 1978). Brackish, estuarine, and inshore waters show less variability in the lead content of sediments; typically, < 10 - 50 $\mu\text{g g}^{-1}$ and up to 850 $\mu\text{g g}^{-1}$ are found in the estuaries of polluted rivers such as the Rhine, or in those with limited water exchange or exposed to the effects of ore mining.

The dumping of sewage sludge may create local hot spots; extreme ranges of 85 - 10 000 $\mu\text{g g}^{-1}$ dry weight have been quoted (Johnson et al., 1974; Nelmes et al., 1974), and more recently 50 - 3000 $\mu\text{g g}^{-1}$ (Förstner & van Lierde, 1979).

The lead content of dredged spoil from harbours, channels, and estuaries has a reported range of 10 - 1450 $\mu\text{g g}^{-1}$, and other town and industrial wastes dumped at sea may also contribute to the lead content (Stanford et al., 1981).

There is much dispute about the fate of lead in estuaries, about how much lead enters the littoral and pelagic biomasses and how much of it is locked away in sediments; this is important because, as a source of lead, anthropogenic input can be several hundred times greater than natural silt input. An example of lead inputs to an estuary in an urbanized and industrialized area in the United Kingdom is given in Table 7 (Murray et al., 1980).

1.4 Transport, transformation, and bioaccumulation

1.4.1 Transport

Atmospheric transport is a major consideration in lead cycling. Various estimates suggest background levels in air, at points remote from man's activities, to be of the order of 0.1 - 1.0 ng m^{-3} (Chow et al., 1969; Murozumi et al., 1969; Egorov et al., 1970; Jernigan et al., 1971), 0.6 ng m^{-3} (Patterson, 1965), with 8 ng m^{-3} (Chow et al., 1972) over uninhabited mountain areas of southern California. Duce et al. (1974) report a gradient over the western North Atlantic ocean of 2 ng m^{-3} offshore increasing to 50 ng m^{-3} towards the eastern shore of the USA. Much higher atmospheric concentrations generally affect coastal seas and waters; for example, Cambray et al. (1975) estimate the total deposition of lead in the North Sea to be 8.1×10^3 tonnes year⁻¹. High inputs are also suggested for Mediterranean coastal bays (Fukai, 1980).

On the basis of measurements of stable and radioactive ²¹⁰Pb isotopes, Schaule &

Table 7. Comparison of lead inputs to the Humber Estuary^a

Rivers	44 kg day ⁻¹	7.3%
Sewage discharges	23 kg day ⁻¹	3.8%
Industrial discharges	9 kg day ⁻¹	1.5%
Sewage sludge dumping	14 kg day ⁻¹	2.3%
Industrial waste dumping	0.3 kg day ⁻¹	0.05%
Dredged spoil	460 kg day ⁻¹	76.3%
Atmospheric input	52 kg day ⁻¹	8.6%
Direct coastal discharges	0.7 kg day ⁻¹	0.15%
Total	603	100%

^a From: Murray et al. (1980).

influx of lead in rivers is much less significant, considering the size of the north Pacific Ocean, than the natural atmospheric input (Fig. 2).

Estimates of global fluxes into the oceans of the northern hemisphere are continually being revised. Chow (1978), using data from Chow & Patterson (1962), suggests a pre-industrial background input of 1.1×10^4 tonnes year⁻¹, supplemented in modern times by a 21×10^4 tonne year⁻¹ input through rain and a 17×10^4 tonne year⁻¹ river input. It is possible that only 1% of riverborne lead is carried beyond the continent shelf. Settlement of lead associated with biological debris (4×10^{10} tonnes year⁻¹ with a lead content of $1 - 10 \mu\text{g g}^{-1}$) is between 4 and 40×10^4 g year⁻¹; much of this is within shelf areas.

1.4.2 Transformation

Atmospheric alkyl-lead contributions fall from high values near petrol handling sites to 1 - 15% of the total, usually in busy cities. It is generally agreed that alkyl leads are degraded to inorganic forms with "a relatively short period" (Nriagu, 1978) in the atmosphere.

There are indications that the survival time of alkyl-lead compounds in surface waters is only a matter of days or weeks and, therefore, values found in sea water or marine organisms ought to be very low. Published high values are suspect (Bernhard, 1980). Although tetra-alkyl leads are the most toxic of the alkyl lead compounds, they are rapidly converted in water to the less toxic tri-alkyl compounds. Biomethylation of lead has been demonstrated in vitro in marine sediments.

1.4.3 Bioaccumulation

A review of several aspects of lead chemobiokinetics has been given by Branica & Konrad (1980) but, in general, little attention has been paid to lead in comparison to other heavy metals. Lead can be taken up, either in the inorganic form or as organolead compounds, from ambient water, sediments, or through diet. The following paragraphs indicate the paucity of information available on these pathways.

Bioavailability of lead discharged in estuaries is discussed by Rickard & Nriagu (1978). Most lead-containing estuarine particulates liberate a major proportion of lead on treatment with weak organic acid. Such indications may be misleading regarding the availability of lead in soluble form, since the normal range of sea water pH would be




Fig. 2. Oceanic outputs, river and atmospheric inputs, and reservoirs of soluble lead at present times in the open North Pacific (Schaule & Patterson, 1980).

Few experiments have been carried out on the toxicity of complexed lead to marine organisms, probably because the concentrations of lead required to produce harmful effects are so high that only a small proportion of lead in natural sea water could be in a complexed form. Canterford & Canterford (1980) estimated that the amount of "free" lead required to reduce by 50% the growth of the algae *Ditylum brightwellii*, in a synthetic sea water medium containing EDTA, was 1.0 - 1.3 $\mu\text{g litre}^{-1}$; comparison with other algae indicates that either this species is abnormally sensitive or that the remaining EDTA-complexed lead was toxic and, thus, by implication, capable of bioaccumulation.

1.4.3.1 Inorganic lead

Bryan (1976) found that *Nereis diversicolor* from sediments containing 8 g lead/kg dry weight contained 1 g kg^{-1} lead dry weight in their tissues. Ray et al. (1981) also found that *N. virens* accumulated lead from sediments containing 243 mg lead kg^{-1} dry weight but not from sediments containing 96 mg kg^{-1} . However, *Macoma balthica* and *Crangon septemspinosa* both accumulated lead from the lesser contaminated sediment. It is not clear whether the lead was absorbed from interstitial water or from sediment particles, although it is most probably the latter, since most of the lead could be extracted by EDTA. Luoma & Bryan (1978) found that the uptake of lead from sediments by *Scrobicularia* was inversely related to the concentration of iron present.

Crustacea

Little is known of the ability of this group of animals to accumulate lead. Weis & Weis (1979) found that the fiddler crab (*Uca pugilator*) exposed to 100 $\mu\text{g lead litre}^{-1}$ in sea water accumulated 2 mg kg^{-1} in 14 days.

Molluscs

Most attention has been given to this group, especially bivalves, and uptake appears to be linear in all species tested. Mussels (*Mytilus edulis*) showed a linear uptake during a 40-day exposure, even at the highest concentration tested, i.e., 5 mg lead litre^{-1} . They accumulated 3.7 g kg^{-1} wet weight in 39 days (Schulz-Baldes, 1974). Similarly, Philipps (1976) exposed mussels to 10 and 20 $\mu\text{g lead litre}^{-1}$ for 35 days and found accumulations of 10.7 and 13.7 mg lead kg^{-1} wet weight, respectively. In these experiments, lead uptake was not affected by zinc, copper, or cadmium in the water. Using a very high ambient concentration of 10 mg lead litre^{-1} , Marshall & Talbot (1979) showed that uptake by the gills of mussels plateaued after 50 days, at which time the gills contained 1 mg lead kg^{-1} .

of 10 weeks (Shuster & Pringle, 1969). Accumulation factors ranged from 1160 to 1400, which was similar to that for mussels.

Young abalone (*Haliotis* sp.) were fed for 6 weeks on brown algae (*Egrecia laevigata*) which had been exposed to 1 mg lead litre⁻¹ and which had accumulated up to 21 mg lead kg⁻¹ wet weight in their tissues (Stewart & Schulz-Baldes 1976). This was not found to have any effect on their growth or activity. Denton & Burden-Jones (1981) found that the Black-lip oyster (*Saccostrea echinata*) accumulated more lead at a salinity of 20 ‰ than at one of 36 ‰, whereas Philipps (1976) found that the lower salinity reduced uptake by the mussel.

In mussels exposed to high ambient concentrations, lead has been found as crystalline extracellular deposits in the capillary walls of the gill (Marshall & Talbot, 1979). These deposits are possibly a mixed or complexed carbonate with calcium. George (1980) reports the presence of lead in membrane-limited vesicles in the gill, and indicates that such granules are excreted by the kidney into the urine. Where half-lives have been measured, these appear to be relatively short. Denton & Burden-Jones (1981) found for Black-lip oysters a half-life of 26 - 34 days which was unaffected by temperature or salinity. Schulz-Baldes (1974) found that mussels lost 33% of accumulated lead with a 40-day exposure to clean water. It is possible that the rate of loss of lead depends, in part, on initial exposure concentrations and on specific detoxification mechanisms.

Echinoderms

Uptake of lead by *Lytechinus pictus* embryos was linear in solutions containing 100 - 1000 µg lead litre⁻¹ (Nash et al., 1981).

Fish

There are few data on the uptake of lead by fish. Somero et al. (1977) exposed an estuarine fish, *Gillichthys mirabilis*, to lead acetate concentrations of up to 2.6 mg lead litre⁻¹. After 100 days, the highest accumulation factors were found in the spleen, gill, and fin tissue, but the factor for muscle tissue was close to unity. Reduction in salinity to 25% sea water doubled the muscle bioaccumulations.

1.4.3.2 Organic lead compounds

These have been little studied in marine organisms. The most comprehensive data have been provided by Maddock & Taylor (1980). The following accumulation factors were found for 3 species exposed to their 96-h LC₅₀ for 4 days (Table 8).

Table 8. Accumulation factors for 3 species exposed to their 96-h LC₅₀ for 4 days

	Shrimp (<i>Crangon crangon</i>)	Mussel (<i>Mytilus edulis</i>)	Plaice (<i>Pleuronectes platessa</i>)
Tetramethyl lead	20	170	60
Tetraethyl lead	650	120	130
Trimethyl lead	1	24	1
Triethyl lead	2	10	2

Further experiments with trimethyl lead showed that a plateau concentration occurred in mussels within about 10 days, and depuration was rapid with a half-life of about 3 days. In contrast, depuration of trimethyl and triethyl lead from dabs (*Limanda limanda*) was very slow, with half-lives in excess of 41 days; with this species, accumulation factors were about 2 and 12 after 41 days' exposure to trimethyl and

Food chains

The transfer factor (concentration in consumer/concentration in prey) for trophic chains in the River Loire (France) was found to be < 1 (Amiard et al., 1980; Amiard-Triquet et al., 1980). So lead does not appear to be accumulated in greater concentrations by top aquatic predators.

However, recent concern has been expressed about 2 potential sources of lead poisoning in birds. There is some evidence that mute swans (Cygnus olor) ingest lead shot or anglers' discarded lead weights which is retained and slowly ground in the gizzard (Birkhead, 1982). Dead swans with lead in their gizzards contained median concentrations of 908 mg lead kg⁻¹ dry weight in the kidney compared with 8 mg kg⁻¹ in dead swans without lead. However, experiments with White Chinese Geese (Johnson & Damron, 1982) fed with lead shot failed to show a lethal effect, and it was proposed that diet may be an important contributing factor.

In the autumn of 1979, over 2000 birds, mainly waders and gulls, were found dead or ill in the Mersey Estuary (United Kingdom); affected birds contained elevated alkyl lead levels; concentration of alkyl lead in the liver of dead dunlin (Calidris alpina) averaged 11 mg kg⁻¹ wet weight. Industrial discharges to the estuary include trialkyl lead, and the mollusc Macoma bathica was found to contain about 1 mg lead kg⁻¹, mostly in the form of alkyl lead. It is thought that the contaminated invertebrate food was the cause of the bird's mortality (Bull et al., 1983).

1.5 Lead in sea water, sediments, and marine biota

1.5.1 Sea water

Since it is clearly important to understand how atmospheric inputs of lead (and other trace metals) interact with the surface microlayer of the sea, attention must be drawn to the recent work of Hunter (1980) on samples collected inshore and in the North Sea. With few exceptions, lead was enriched in all microlayer samples compared to sub-surface concentrations, probably as a result of flotation of particles attached to rising bubbles and perhaps assisted by direct influx of atmospheric particles.

According to Burnett & Patterson (1980), the shallow waters of the open (Pacific) ocean contain only 10 ng lead kg⁻¹ and deep ocean waters, 1 or 2 ng lead kg⁻¹. The range of values for the North Atlantic Ocean and around the British Isles is large and, 50 - 1200 ng litre⁻¹ (Topping et al., 1980).

The generalized depth profile for lead in the northeast Pacific Ocean (Fig. 3(a)) closely resembles those for ²¹⁰Pb and tritium (Fig. 3(b)). These figures are interpreted by Schaule & Patterson (1980) as demonstrating the contemporary incorporation of airborne lead mainly from lead in petrol simultaneously with ²¹⁰Pb from continental ²²²Rn emanations and tritium from nuclear bomb test debris.

Earlier estimates of the residence time for lead in the ocean were of the order of 10³ years (Chow, 1978). Recently, Schaule & Patterson (1980) have lowered the estimate to only 2 years in the surface layer, and about 200 - 400 years in deep water.

1.5.2 Sediments

From high values in certain estuaries, the lead content of open sea sediments, 8.4 - 60 µg g⁻¹ (Nriagu, 1978), falls off away from the coast but retains considerable spatial heterogeneity. Deep-sea sediments have ranges of 13 - 17 µg g⁻¹ for oozes, 47 - 61 µg g⁻¹ for clays, and 740 - 1250 µg g⁻¹ for manganese nodules. Local enhancement is reported for areas influenced by hot brine deposits and volcanic activity.

Several reports confirm the recent enhanced deposition of lead in sediment cores from ponds, lakes, and the marine environment from many areas. One example (Fig. 4) is quoted from Bertine (1980) which considers a variety of depositional areas.

1.5.3 Marine biota

Data on the lead content of marine biota are given in Table 1 (1980).




Fig. 3. (a) Generalized depth profile of common lead concentration in the central northeast Pacific Ocean.
 (b) General structure of tritium and ^{210}Pb distribution in the central northeast Pacific Ocean.
 From: Schaule & Patterson (1980).




Fig. 4. Lead content (\bullet) and lead/aluminium ratios (\circ) in Southern California Basin sediments, and Whites Point sewer outfall sediments; note change of scale, Whites Point. Vertical dating of sediments is indicated. From: Bertine (1980).

due to sample contamination. Similar reserves have to be made for the extensive surveys of trace metal concentrations in commercial fish and shellfish species coordinated by ICES in the northern Atlantic, although the participating laboratories intercalibrated (ICES, 1974, 1977a,b,c, 1980). Most of these concentrations reported are below the detectable limit of $0.01 \mu\text{g g}^{-1}$ fresh weight, and probably the only conclusion that may be drawn from these data is that the lead levels cannot be higher. The very low values of Patterson & collaborators, which may have been accepted generally as true levels, because they were carried out under ultra-clean conditions (Patterson & Seattle,

1976; Burnett & Patterson, 1980) may serve as a guideline for the concentrations to be expected:

Species	µg/g fresh weight
Valonia, alga	0.2
Abalone, muscle	0.004
<u>Mytilus</u> , muscle	0.025
Spiny lobster, muscle	0.005
Tuna, muscle	0.0003

Davies (1981) and Murray (1979) provide in-depth data from routine surveys of northern and southern North Sea commercial fish and shellfish. The coverage is unequal with regard to the components of the food web for representative sea areas and, for the most part, it is inadequate when defining the contributions made by species, age, sex, feeding habits, maturation, exposure to natural or anthropogenic lead levels, and so on. In particular, the data on lead in phytoplankton and its members, which is critical in defining the role of primary production on the fate of lead in the sea, is almost totally lacking because of uncertainties in sampling and contamination.

Phytoplankton

Using Eisler (1983) data, except where otherwise indicated, the only guideline value for lead in phytoplankton is 40 µg g⁻¹ dry weight.

Macrophytes

A selection of macrophytes, including edible seaweeds, exhibit accumulation factors from 1200 to 82 000. Red, green, and brown seaweeds commonly have lead contents of 3.0 - 20 µg g⁻¹ dry weight. Lead contents can reach a maximum of 300 µg g⁻¹ dry weight depending on species and ambient lead concentrations under either natural or anthropogenic influences. Sea grasses are important food for certain wild fowl, and their typical lead concentrations (µg g⁻¹) are 5.0 for *Spartina*, 1.6 - 8.4 for *Zostera*, and up to 1800 (all dry weight) for *Zostera* (entire).

Zooplankton

Values given for pelagic zooplankton are: anomalocera 3.5 - 6.0, copepods 3.3, meganyctiphanes 2.2, and euphausiids 2.1 µg g⁻¹ dry weight. All suggest moderate lead accumulation. Pelagic carnivores such as *Aurelia* sp. (0.8 µg g⁻¹ dry weight), *Beröe* sp. (6.0 µg g⁻¹ ash weight), and *Cyanea* sp. (6.0 µg g⁻¹ ash weight) have low lead contents.

Crustacea

Larger filter feeders such as shrimp (0.2 - 1.2 µg g⁻¹ wet weight) and rock shrimp (1.6 µg g⁻¹ dry weight) also have low lead contents in their soft parts, with somewhat higher levels in the exoskeleton. The important benthic crustacean, *Nephrops norvegica* (< 0.1 - 0.5 µg g⁻¹ wet weight), *Cancer pagurus* (< 0.1 - 0.3 µg g⁻¹ wet weight), and *Homarus vulgaris* (0.4 µg g⁻¹ wet weight) (Davies, 1981) sampled from commercial fishing areas all have very low lead contents.

Molluscs

Common benthic animals, such as corals with 2 - 42 µg g⁻¹ dry weight show response to depth; *Alcyonium* (24 µg g⁻¹ weight), abalone, cardium, chlamys, oysters, littorina, modiolus, and *Mya* sp. are all well below 5 µg g⁻¹ wet weight. Several species, at least, show high values in specific organs, e.g., *Chlamys* kidney, 830 µg g⁻¹ dry weight; *Pecten* kidney, 137 000 µg g⁻¹ dry weight. For *Mytilus* sp., used in the Mussel Watch Programme (Goldberg et al., 1978) very many lead values are

harvested areas, have lead contents not exceeding $1 \mu\text{g g}^{-1}$ wet weight. In the Eisler (1983) compilation, values up to $450 \mu\text{g g}^{-1}$ dry weight are reported. Accumulation factors for soft parts range from thousands to tens of thousands. Accumulation is variable in the shell but, in gills and visceral tissues, it is usually considerable. Some squids and octopuses which are also commercially fished have values of $< 0.4 \mu\text{g g}^{-1}$ wet weight.

Fish

The lead contents of all common commercial fish species sampled in the United Kingdom are consistently low and typically lower than the detection limit of approximately $0.1 \mu\text{g g}^{-1}$ wet weight (Murray, 1979; Davies, 1981). Diverse observations on elasmobranchs and many teleost species give most values at $< 0.5 \mu\text{g g}^{-1}$ wet weight, and extreme values which seldom exceed $1.0 \mu\text{g g}^{-1}$ wet weight. Exceptions are Ciliata sp. at $8.1 - 25 \mu\text{g g}^{-1}$ dry weight, Chinocottus sp. at $0.6 - 4.9 \mu\text{g g}^{-1}$ wet weight, "flounder meal" with total lead at $5.3 \mu\text{g g}^{-1}$ and tetraalkyl lead at $4.8 \mu\text{g g}^{-1}$ wet weight, and Platichthys sp. at $14 - 28 \mu\text{g g}^{-1}$ dry weight. The lead content of flesh (muscle) is usually lower than that of skin and internal organs, but the disproportion is minor, unlike in molluscs, and the response to environmental lead is small for most edible species (perhaps with the exception of flounder).

Birds

In the kidney and liver of sea birds, lead contents are modest, namely < 2.1 and $< 5.3 \mu\text{g g}^{-1}$ wet weight, but information is meagre.

Mammals

Fish-eating mammals (4 spp) show very low values except for the harbour seal (Phoca). It has a lead accumulation in the kidney ranging from $0.08 - 0.60$ and in the liver from $0.09 - 5.3 \mu\text{g g}^{-1}$ wet weight. Higher levels are common in the hard tissues.

2. Effects on Marine Biota

2.1 Reference documentation

A summary of the information on acute and chronic effects of lead on marine organisms has been published by the US EPA (1980), and tabular summaries listing aqueous concentrations and their effects on organisms have been prepared by Taylor (1981) and IRPTC (1981). Individual research papers are listed in the reference section, but the following critical review is based mainly on those studies in which low concentrations have been shown to have an effect.

2.2 Effects on marine biota

2.2.1 Inorganic lead

Algae

Although most of the published data indicate that concentrations greater than $100 \mu\text{g lead litre}^{-1}$ are required to produce acute or chronic effects on algae, there are a number of papers which report effects at lower levels.

Using a natural population of mixed algal species in enriched sea water, Hollibaugh et al. (1980) found that Chaetoceros sp. were affected by $60 \mu\text{g lead litre}^{-1}$. Other algae increased in numbers, and a slight growth reduction of Thalassira aestivalis occurred at $100 \mu\text{g litre}^{-1}$. Similar results were obtained for Phaeodactylum tricornutum (Woolery & Lewin, 1976) in that $100 \mu\text{g lead litre}^{-1}$ (as PbCl_2) reduced photosynthesis (but not respiration) to 70 - 80% of control values. Using synthetic sea water, Rivkin (1979) found that $4.4 - 7.8 \mu\text{g lead litre}^{-1}$ caused a 50% reduction in chlorophyll and cell numbers (compared with controls) of Skeletonema

EDTA-complexed lead in synthetic sea water, Canterford & Canterford (1980) estimated that only 1.0 - 1.3 μg "free" lead litre⁻¹ was required to reduce growth of Ditylum brightwellii to 50% of control values. The significance of these low effect levels in synthetic sea waters is unclear.

Protozoa

The growth rate of Cristigera was slightly reduced by 300 μg lead litre⁻¹ (Gray & Ventilla, 1973).

Annelida

This group of organisms appears to be resistant to lead toxicity. Juvenile polychaetes (Neanthes arenacoedentata) exposed to dilutions of lead citrate in sea water had a 28-day LC₅₀ of 2.5 mg lead litre⁻¹, which compared with 3.2 mg litre⁻¹ for adults of the same species (Reish et al., 1976). These authors also found that the 96-h LC₅₀ for trochophore larvae of Capitella capitata was 1.2 mg lead litre⁻¹, which compared with a 28-day LC₅₀ of 1.0 mg litre⁻¹ for the adult. Similar experiments by Reish & Carr (1978) on Ctenodrilus serratus showed that the 96-h LC₅₀ was > 20 mg lead litre⁻¹, and that concentrations > 1.0 mg litre⁻¹ reduced population sizes over a 21-day exposure period. The population size of Ophryotrocha diadema (96-h LC₅₀ of 11 mg lead litre⁻¹) was also reduced by lead concentrations in the range 1 - 5 mg litre⁻¹. Brown & Ahsanullah (1971) found that 1.0 mg lead litre⁻¹ killed less than 80% of a batch of Ophryotrocha labronica in 25 days, and 10 mg litre⁻¹ did not suppress growth within an 8-day exposure period. Therefore, this group of organisms appears to be resistant to lead.

Crustacea

These organisms, too, appear to be resistant to lead toxicity. Embryonic development of the mud crab Rhithropanopeus harrisi to the megalopa stage was delayed from 14.3 days to 15.4 days by 50 μg lead litre⁻¹ (Benijts-Claus & Benijts, 1975); the biological significance of such a short delay is unclear. Martin et al. (1981) found that the 48-h LC₅₀ for Cancer magister zoea was 575 μg lead litre⁻¹. Chaisemartin et al. (1978) found increased production of aspartic amino-transferase in the hepatopancreas of the crab Macropodia rostrata exposed to 102 μg lead litre⁻¹. However, Zencirci (1980) found that 100 μg lead litre⁻¹ produced 100% mortality of Gammarus locusta in 18 days. The 24-day LC₅₀ for Artemia salina was found to be 1.0 mg lead litre⁻¹, but this concentration did not reduce growth rates within a 10-day exposure period (Brown & Ahsanullah, 1971).

Mollusca

Juvenile stages of molluscs are usually more sensitive to toxic substances than are adults. However, Calabrese et al. (1973) found that the no-observed-effect level for oyster (Crassostrea virginica) embryos was 500 μg litre⁻¹ for a 48-h exposure period. Similar tests with C. gigas and Mytilus edulis embryos gave 48-h EC₅₀s of 758 and 476 μg lead litre⁻¹, respectively (Martin et al., 1981), 760 μg lead litre⁻¹ for C. virginica, (unpublished MAFF data), and 780 μg lead litre⁻¹ with a no-effect level of 400 μg litre⁻¹ for Mercenaria mercenaria (Calabrese & Nelson, 1974). Using Mytilus galloprovincialis, Hrs-Brenko et al. (1977) found that embryo development in lead solutions was affected by an increase in temperature and a reduction in salinity. With a salinity of < 32.5 ‰, abnormalities increased at > 17.5 °C and 100 μg lead litre⁻¹ and at 15 °C with 250 μg lead litre⁻¹.

Lethal concentrations reported for juvenile and adult molluscs are in excess of 1 mg lead litre⁻¹. Exposure of Macoma balthica to 500 μg lead litre⁻¹ for 24 h at a salinity of 6 ‰, and 6 °C slightly reduced burrowing activity when they were returned to clean water; a concentration of > 5 mg litre⁻¹ was required to damage the siphons (Eldon et al., 1980). Data on the long-term accumulation of lead by molluscs, reviewed in a previous section, provides additional evidence of resistance of these species with prolonged exposure.

Echinoderms

Tests with developing sea urchin embryos over a 12-h exposure period indicated lead concentrations of between 1.1 and 2.2 mg lead litre⁻¹ (added as lead acetate) at a maximum no-observed-adverse effect on eggs of Anthocidaris crassispina to reach the gastrula stage (Kobayashi, 1971). Retardation of plutei development of Arbacia punctulata was observed at 10 mg lead litre⁻¹ (or PbCl₂) over a 13.5-h exposure period (Waterman, 1937), and the true exposure concentration was probably lower.

Fish

Few experiments have been carried out with marine fish. In Fundulus heteroclitus, exposure to 1 mg lead litre⁻¹ did not retard fin regeneration within a 14-day exposure period (Weis & Weis, 1979). When mullet Mugil auratus were exposed to about 470 µg lead litre⁻¹, the blood lead levels rose linearly for 28 days, whereas a decrease in erythrocyte ALA-D and haemoglobin content plateaued within this period (Krajnovic-Ozretic & Ozretic, 1980). After 80 days exposure, blood haemoglobin fell from 8.8 to 5.5 g 100 ml⁻¹, and ALA-D activity decreased by about 50%.

2.2.2 Organic lead

The toxicity of the various alkyl lead compounds are summarized in Table 9.

Table. 9. The toxicity of various alkyl-lead compounds

	Maddock and Taylor (1980)				Marchetti (1978)		
	96-h LC ₅₀ (mg litre ⁻¹)			6-h EC ₅₀ (mg litre ⁻¹)	48-h LC ₅₀ (mg litre ⁻¹)	48-h EC ₅₀ (mg litre ⁻¹)	
	Shrimp	Mussel	Plaice	Alga ^a	Artemia	Morone labrax	Alga ^b
Tetra-methyl lead	0.11	0.27	0.05	1.3	0.25	0.10	1.65
Tetra-ethyl lead	0.02	0.10	0.23	0.1	0.085	0.065	0.15
Tri-methyl PbCl	8.8	0.5	24.6	0.8			
Tri-ethyl PbCl	5.8	1.1	1.7	0.1			
Dimethyl PbCl ₂			300				
Diethyl PbCl ₂			75				

^a Phaeodactylum tricorutum; 50% reduction in photosynthetic activity.

^b Dunaliella tertiolect; 50% reduction in photosynthetic activity.

Although tetra-alkyl leads are the most toxic form of lead, they are rapidly converted to the less toxic tri-alkyl compounds in water. There appears to be agreement between the 2 sets of data. Although the studies on tetra-alkyl lead by Maddock & Taylor (1980) were under continuous flow conditions of test (the remaining ones being static), those of Marchetti (1978) were static, but in closed vessels. Although the toxicity of the tetra-alkyl leads was greater than that of the inorganic forms, the bioconcentration factors were much lower. Tests with oyster larvae (unpublished MAFF data) showed that the 48-h EC₅₀ for tri-methyl lead was 0.1 mg lead litre⁻¹.

3. Human Health Aspects

3.1 Reference documentation

Several reviews on health effects of lead have been published (US EPA, 1977; WHO, 1977; Jaworski, 1979; Bornschein et al., 1980; DHSS, 1980; Needleman, 1980; Rutter, 1980; WHO, 1980; Ratcliffe, 1981; Chisolm & O'Hara, 1982; Oskarsson & Camner, 1983; Rutter & Russel-Jones, 1983). Due to the great number of recent scientific reports not included in the reviews, the following section is primarily based on original publications which are listed in the reference section.

3.2 Toxicokinetic properties

The absorption of lead from the gastrointestinal tract varies considerably. Balance studies in human beings have shown that about 10% of the lead in food is absorbed through the gastrointestinal tract (WHO, 1977b). In addition to individual variation, the absorption is highly dependent on the presence of food in the gastrointestinal tract. Flanagan et al. (1982) studied the retention of radioactive lead in 85 fasting subjects and found that approximately 60% of an oral dose of 4 - 400 μg of lead was retained. The effect of minerals on uptake of lead in the gastrointestinal tract was studied by Heard & Chamberlain (1982), who found that addition of calcium and phosphate in doses equivalent to that which a normal meal contains reduced the uptake of lead from 60 to 10%. The chemical species of lead have some influence on the absorption, and water soluble lead compounds are more easily absorbed than those which are less soluble (Chamberlain et al., 1978).

The effect of age on gastrointestinal absorption of lead has been studied in experimental animals. Kostial et al. (1971) demonstrated that 5 - 7 day-old rats absorb about 55% of a single oral dose of radioactive lead. There are very few data on the gastrointestinal absorption of lead in children. Ziegler et al. (1978) performed 89 balance studies with 12 healthy infants ranging in age from 14 days to 2 years. With lead intakes above 5 $\mu\text{g kg}^{-1} \text{ day}^{-1}$, they reported a mean absorption of 41.5% with large individual variations. Eleven lead balance studies were performed on 8 healthy children ranging in age from 3 months to 8.5 years (Alexander et al., 1973, 1974). An absorption of 53% was reported, and the mean retention, calculated as intake minus total excretion, was 18% with a range of -4% to 37%. The small number of measurements and the large range of values do not allow any conclusions to be drawn about the relation between gastrointestinal absorption of lead and age in children.

The body burden of lead is divided into 2 fractions: one firmly bound to bone and another loosely bound to blood and soft tissues. The bone fraction constitutes about 90% of the total body burden of lead. The mean retention of lead in blood and soft tissues is about 3 weeks to 1 month and, in bone, about 5 years (Chamberlain et al., 1978; Schütz et al., 1981).

Lead is excreted via urine and faeces. The faecal excretion is mainly unabsorbed lead and could be used as an indicator of the oral intake of lead.

Lead in blood (PbB) is a good indicator of current lead exposure. PbB can be related to the intake of lead, the concentrations in air, diet, or water, and to different health effects due to lead exposure.

PbB levels vary in different parts of the world. A study was performed on PbB levels in the populations of big cities in member countries of the European Community (EEC) (Berlin, 1982). UNEP/WHO initiated a global study in which the PbB levels in teachers, living in big cities in different parts of the world, were measured (Vahter, 1982). The analyses were made with emphasis on quality control in order to obtain accurate and intercomparable results. Figs. 5 and 6 summarize the results from the 2 studies. The lowest PbB levels were found in Tokyo, Peking, Stockholm, and Baltimore, where the median levels were below 10 $\mu\text{g}/100 \text{ ml}$. The US national estimates of PbB levels from 1976 to 1980 have been reported recently (Mahaffey et al., 1982). Blood samples were analysed from a total of 27 800 persons, representing age groups from 6 months to 74 years of age. The results show that 22% of the whole population had PbB levels under 10 $\mu\text{g}/100 \text{ ml}$ and 1.9% had levels above 30 $\mu\text{g}/100 \text{ ml}$.

Among children in Sweden, Finland, and Denmark, the PbB levels are about the same as or a little lower than in adults. The mean PbB levels in children living in big cities are about 10 $\mu\text{g}/100 \text{ ml}$.




Fig. 5. Lead levels in blood (median and 90-percentile in the population of European cities. From: Berlin (1982).




Fig. 6. Lead levels in blood in teachers (Stockholm, a randomly-selected population) (median and 90-percentile). From: Vahter (1982).

The lowest reported PbB levels are from a population living in a remote mountain area in Nepal (Piomelli et al., 1980). The concentration of lead in air was below the detection limit of $0.004 \mu\text{g m}^{-3}$. The mean value of PbB was $3.4 \mu\text{g}/100 \text{ ml}$. A study on PbB levels in children living in rural areas of Papua New Guinea reported a mean value of $5.2 \mu\text{g}/100 \text{ ml}$ (Poole et al., 1980).

A study in Sweden showed no significant difference in PbB levels in persons living in city areas with high traffic density compared to persons from low traffic areas (Elinder et al., 1983). Tobacco smoking and alcohol consumption have been demonstrated to increase the PbB levels (Olsen et al., 1981; Shaper et al., 1982; Elinder et al., 1983).

It is of great importance to determine the relationship between dietary lead intake and PbB. Chamberlain et al. (1978), using the same model that had been used for inhalation of lead, predicted an increase in PbB of $3.6 \mu\text{g}/100 \text{ ml}$ per $100 \mu\text{g day}^{-1}$ of lead in the diet. The calculations are based on a 15% gastrointestinal absorption, a 50% distribution into a blood volume of 5.4 litre, and an 18-day half-life of lead in blood. The results from 4 human experimental studies cited by Chamberlain showed a range of $1.4 - 4.3 \mu\text{g}/100 \text{ ml}$ per $100 \mu\text{g day}^{-1}$ of lead in the diet. However, this calculation does not seem to be generally valid for estimating the resulting PbB level from a certain intake of lead. A low daily intake of lead of, for example, $30 \mu\text{g day}^{-1}$, as in Sweden, will give too low a PbB value of about $1 \mu\text{g}/100 \text{ ml}$ compared to the measured value of approximately $8 \mu\text{g}/100 \text{ ml}$. If it can be assumed that the intake rate has been correctly determined, the discrepancy can be explained by a non-linear relationship between ingested lead and PbB. The value predicted by Chamberlain et al. (1978) is valid for the determination of the increase of PbB after addition of lead to the diet in the high exposure range. Curvilinearity means that, as the dose increases, the rise in PbB becomes progressively smaller. Such a relationship has been demonstrated by Moore et al. (1977) between lead in drinking-water and PbB, and by Hammond et al. (1981) for the relationship between lead in air and PbB. According to the DHSS (1980), using the equation from Moore et al. (1977), a small increase of lead by ingestion increases PbB by about $7 \mu\text{g}/100 \text{ ml}$ per $100 \mu\text{g day}^{-1}$ of ingested lead when starting from a baseline PbB of $18 \mu\text{g}/100 \text{ ml}$. This becomes $2 \mu\text{g}/100 \text{ ml}$ per $100 \mu\text{g day}^{-1}$ when starting from a baseline of $24 \mu\text{g}/100 \text{ ml}$, and it is about $1 \mu\text{g}/100 \text{ ml}$ per $100 \mu\text{g day}^{-1}$ when starting from a baseline of $30 \mu\text{g}/100 \text{ ml}$. The relationship between first draw water which is in the equation and the ingestion rate of lead was, however, not specified.

3.3 Health effects

Early biochemical changes due to elevated lead exposure occur in the haematopoietic tissues where the biosynthesis of haem is disturbed. The haem biosynthesis pathway and the effect of lead in different steps are summarized in Fig. 7. The most sensitive effect is the inhibition of ALA-D, which has been observed at PbB levels above $10 \mu\text{g}/100 \text{ ml}$ (Nordman, 1975). The decreased activity of ALA-D results in increased urinary excretion of ALA, and this has been reported at $40 \mu\text{g}/100 \text{ ml}$ blood (Zielhuis, 1975). Increased levels of free protoporphyrins in the erythrocytes (FEP) have been associated with PbB levels of about $20 - 30 \mu\text{g}/100 \text{ ml}$ in adult females, about $25 - 35 \mu\text{g}/100 \text{ ml}$ in adult males, and about $15 \mu\text{g}/100 \text{ ml}$ in children (Roels et al., 1975; Zielhuis, 1975; Piomelli et al., 1982). These changes in haem biosynthesis are usually not considered to be adverse health effects as such, but they can be taken as indicators of biological response to elevated lead absorption (US EPA, 1977). At higher PbB levels, there is an effect on the overall haemoglobin synthesis and anaemia will result. Lead-exposed workers developed anaemia at PbB levels of $60 - 80 \mu\text{g}/100 \text{ ml}$ (Baker et al., 1979). Children are more sensitive and anaemia has been reported at PbB levels above $40 \mu\text{g}/100 \text{ ml}$ (Betts et al., 1973).

Lead exposure may have serious effects on the central and peripheral nervous systems. CNS effects are most frequent in children, and PNS effects occur after long-term exposure in adults. PNS defects range from paresis to slight functional impairment. The major neurophysiological disturbances consist of slowing of the motor conduction velocity (especially of the slower fibres), slowing of the sensory conduction velocity, and electromyographic disturbances (WHO, 1980). Reduced conduction velocities begin to occur in the PbB range $40 - 50 \mu\text{g}/100 \text{ ml}$ and become more prominent in the




Fig. 7. Haem biosynthesis and the effect of lead.

Lead in high doses causes encephalopathy. The symptoms are ataxia, coma, and convulsions. In children, lead encephalopathy has occurred at PbB levels above 60 µg/100 ml and, in adults, above 80 µg/100 ml (WHO, 1977b). Neurological sequelae can follow in severe or repeated episodes of lead encephalopathy. The sequelae are commonly of a subtle nature involving impaired learning ability, motor incoordination, disturbed sensory perception, and inability to concentrate (WHO, 1977b). Such disturbances are suspected to follow lower-dose lead exposure than those which cause encephalopathy. The major concern today is the sub-clinical effects of lead on the developing CNS. Effects of lead exposure on intelligence and behaviour has been the subject of many epidemiological studies.

Reports in the older literature suggested that occupationally lead-exposed women had a higher frequency of stillbirths and miscarriages than normal (Rom, 1976). PbB levels were not reported, but these women were probably exposed to very high levels of lead. From animal experiments, it is known that lead passes the placental barrier and can cause CNS damage to the exposed fetus. What effects lower levels of lead exposure have on reproduction in human beings are not known, but it is suspected that lead exerts toxic effects on the developing CNS of the fetus.

The literature concerning the chromosomal abnormalities in persons exposed to lead is controversial. Most studies have not been able to exclude the possibilities of

exposure. Animal studies have shown that some lead compounds induce renal tumours after peroral administration. In epidemiological studies, it has not been confirmed that lead is also a human carcinogen. Due to the lack of adequate human data, IARC (1980) recommended that, based on experimental animal data, lead acetate, lead subacetate, and lead phosphate be considered as carcinogens.

The major concern in studying health hazards resulting from low-level lead exposure is the possibility that subtle effects on the developing nervous system may result from absorption of lead in smaller quantities than those known to give symptoms. There have been a number of studies conducted on children exposed to lead where intelligence and behaviour have been measured. There are methodological problems with these studies due to imprecise measurements of the intensity and duration of the lead exposure, varying definitions of the nature of the insult, and several confounding variables which are hard to control. Several critical reviews of the literature on this subject have been published lately (Bornschein et al., 1980; DHSS, 1980; Needleman, 1980; Rutter, 1980; Ratcliffe, 1981; Rutter & Russel-Jones, 1983).

The most common indicator of lead exposure is PbB. A single blood sample gives an indication of the present lead exposure but does not show the exposure in earlier childhood. A normal PbB level does not rule out previous chronic lead exposure, and an elevated PbB value may reflect an occasional high exposure. Repeated measures of PbB levels over a long time-span would be preferable. Determination of lead levels in dentine could be a useful measure of past exposure. There is a moderate but not high correlation between dentine and blood lead levels.

A battery of tests for measures of intelligence, general cognitive development and behaviour, as well as some for neurologic and psychomotor functions, have been used. Diverse tests have been used in different studies, and the results are hard to compare.

In all probability, there are disparities between test and control groups other than PbB levels which may affect the performance of the neurological and behavioural experiments. Confounding variables such as age, sex, socio-economic status, and parental IQ should be controlled. There are other variables more difficult to measure which also affect development and behaviour, such as the quality of the care-giving environment (Milar et al., 1980).

Taking all evidence into consideration from recent animal as well as human studies (US EPA, 1977; WHO, 1977; Needleman et al., 1979; Ernhart et al., 1981; Otto et al., 1981; Yule et al., 1981; Rice, 1982; Silbergeld, 1982; Lansdown et al., 1983; Winneke et al., 1983), it appears that severe encephalopathy has occurred at around 60 $\mu\text{g}/100\text{ ml}$ in blood. Minor neuropsychological effects may occur at considerably lower blood lead levels (around 35 $\mu\text{g}/100\text{ ml}$), and there are some indications that effects may occur in sensitive groups already at levels of 15 - 30 $\mu\text{g}/100\text{ ml}$. Scientific consensus on effects at these low levels has, however, not yet been reached.

3.4 Total exposure to lead

Of the 3 routes of exposure, food, water and air, food is considered the major contributor to the total lead exposure (NF, 1982). Lead in food can be derived naturally or as a result of man's activities from direct (e.g., dustfall on crops) or indirect contamination (e.g., dustfall on soil, contaminated water for irrigation, or sewage sludge used as fertiliser). Furthermore, lead can get into food through food processing. Lead-soldered cans for food packaging have been shown to increase the lead content in the food to a great extent (DHSS, 1980; NF, 1982).

The DHSS (1980) estimated the contribution of lead from different sources to the body burden. Based on the mean blood lead and the contribution from air as described in section 3.3.4.2, the percentage of blood lead from food varied between 44% and 91%, the lower percentage calculated from an air lead level of 1 $\mu\text{g m}^{-3}$ and a water lead level of 50 $\mu\text{g}/\text{litre}$ (DHSS, 1980). These calculations are, of course, approximate. Schaffner (1981), citing Mahaffey (1981), estimated that 55 - 85% of a person's daily exposure originates from food. Tsuchiya (1979) gave an estimation of 80 - 85%.

3.4.1 Dietary lead intake

Daily dietary intakes of lead in different countries and age groups are shown in Table 10. There are 2 ways of estimating the lead intake from food (DHSS, 1980). One way is to calculate the lead content in foods made up to represent an average national diet (total diet studies), the other is to analyse duplicate portions of food actually consumed (individual diet studies). The mean daily dietary intake of lead calculated from total diets was 113 μg in Great Britain compared to 75 μg when calculated from individual diets (Table 10). There are many factors to consider when estimating the dietary intake of lead. For example, differences in the composition of diet in various parts of the world could account for differing intakes of lead. In some cases, the variations among countries might partly be explained by methodological differences. Very seldom has the analytical quality been controlled and, therefore, reliable comparisons between different studies can hardly be made.

Table 10. Estimated average daily intakes of lead ($\mu\text{g person}^{-1} \text{ day}^{-1}$)

Country	Age, sex	Lead intake ($\mu\text{g/day}$)	Reference
Canada	0 - 5 months	33	NF (1982)
Canada	40 - 64 years; male	113	NF (1982)
Canada	40 - 64 years; female	89	NF (1982)
Finland	adults	66	Varo & Koivistoinen (1980)
Italy	adults	400	Tomassi, National Institute of Nutrition Italy (personal communication)
Sweden	adult; male	27	Slorach et al. (1982)
Sweden	50 - 60 years; male	33	Schütz (1979)
United Kingdom	0 - 4 months	17	DHSS (1980)
United Kingdom	adults ^a	113	DHSS (1980)
United Kingdom	adults ^b	75	DHSS (1980)
USA	0 - 5 months	20 - 46	NF (1982)
USA	2 - 6 years	60 - 70	NF (1982)
USA	teenage; male	79 - 95	Jelinek (1982)

^a Calculated from total diets.

^b Calculated from individual diets.

A provisional, tolerable weekly intake of 3.0 mg of lead for adults was recommended by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) in 1972 (FAO/WHO, 1972). This is equivalent to about 430 $\mu\text{g day}^{-1}$. The recommendation does not apply to children and any increase in the amount of lead derived from drinking-water or inhaled from the atmosphere will reduce the amount tolerated through food intake. What PbB level such an intake corresponds to is unknown and, therefore, it is not possible to relate this intake level to the health effects of lead.

Total dietary lead intake in children is lower than in adults. Lead intakes in $\mu\text{g day}^{-1}$ increase with age, but on body weight basis ($\mu\text{g kg}^{-1} \text{ day}^{-1}$) decreases with age. Thus, for one-year-old children, the mean lead intake was 61 $\mu\text{g day}^{-1}$ or 5.4 $\mu\text{g kg}^{-1} \text{ day}^{-1}$, while for adults 40 - 64 years old, the mean lead intake was 113 $\mu\text{g day}^{-1}$ (males) or 1.6 $\mu\text{g kg}^{-1} \text{ day}^{-1}$ (NF, 1982). A factor of great importance in the comparison of dietary lead intake is the proportion of canned foods included in the diet.

Lead-soldered tin cans constitute a major source of lead in foods. Acidic foods,

inclusion of canned foods in the studied diets will considerably increase the dietary intake. Schütz (1979), in examining dietary intake of lead in Sweden, found 2 diets containing canned fruits that contributed 157 and 167 $\mu\text{g day}^{-1}$ compared to the average 33 $\mu\text{g day}^{-1}$. Slorach et al. (1982) estimated dietary intake of lead in Sweden to be 27 $\mu\text{g day}^{-1}$. Very few foods from lead-soldered cans were included in their study. Jorhem & Slorach (1979) compared lead levels in canned and fresh fruits and vegetables and found a 6- to 28-fold higher level in the canned products. For children 0 - 6 years old, lead from canned foods contributed 17 - 28% of the mean total dietary lead intake (NF, 1982). In the United Kingdom, lead-soldered cans are estimated to contribute approximately 15% of the total dietary intake of lead (DHSS, 1980).

Table 11 summarizes some of the data on lead levels in foods and compares lead levels in fresh and canned food (Jelinek, 1982; NF, 1982). Canned foods contain markedly higher lead levels than fresh foods. This is most evident in fruits.

Because of the toxicity of lead to the developing nervous system, there has been great concern over the need to reduce the levels of lead in foods for infants. In the USA, lead levels in such foods have been reduced 5 - 10 times since the early 1970s, mostly due to the introduction of steel cans or glass containers instead of welded cans (Jelinek, 1982).

Table 11. Lead content in foods ($\mu\text{g g}^{-1}$ fresh weight)^a

Food	Uncanned	Canned
Dairy products and eggs		
Milk	0.02	0.10 - 0.13
Butter	0.07	
Ice cream	0.01	
Cheese	0.05	
Eggs	0.17	
Meat and poultry		
Beef, pork, lamb, veal	0.06	0.24
Poultry	0.12	0.24
Hamburger	0.25	
Beef liver	0.09	
Cereal, nut and sugar products		
Flour, white	0.05	
Bread, white	0.08	
Cereals, breakfast	0.11	
Peanut butter	0.06	
Sugar, refined	0.03	
Vegetables		
Potatoes	0.05	0.12
Cabbage	0.01 - 0.04	0.08
Lettuce	0.12 - 0.15	0.39
Beans	0.01 - 0.04	0.16 - 0.32
Peas	0.03	0.27
Carrots	0.14	0.13
Onions	0.18	0.32
Tomatoes	0.05 - 0.08	0.30 - 0.37
Cucumbers	0.02	

Table 11 (contd).

Fruits		
Citrus (oranges, lemons)	0.01	0.39
Apples	0.02	0.22
Cherries	0.02	0.39
Pears	0.02	0.18 - 0.19
Fish and shellfish		
Salmon	0.39	0.72
Mackerel	0.40	0.99
Tuna		0.45
Cod	0.06	
Flounder	0.10	
Oysters	0.17	
Clams	0.21	

^a From: NFPA/CMI Report for the US FDA (1980) cited in NF (1982); Jelinek (1982).

Lead levels in Swedish human breast milk have been analysed in a study initiated by WHO (Larsson et al., 1981). The median lead content in 41 samples was $0.002 \mu\text{g g}^{-1}$ with a range of $0.0005 - 0.009 \mu\text{g g}^{-1}$. The calculated daily intake for a 3-month-old infant who consumed only breast milk was $0.27 \mu\text{g kg}^{-1}$ body-weight.

Lead intake from wine can be an important contribution to the daily intake of heavy wine consumers. Noirfalise & Collinge (1982) analysed the lead content in samples of French and Italian wines and reported concentrations in the range $18 \mu\text{g}^{-1}$ to $262 \mu\text{g litre}^{-1}$. The average concentration was $65 \mu\text{g litre}^{-1}$.

3.4.2 Exposure from water and air

Lead from tap water not only contaminates food during cooking but also contributes to lead intake through drinking. Normally, tap water contains less than $10 \mu\text{g litre}^{-1}$, but markedly higher concentrations are found where lead piping carries soft water. In Great Britain, 10% of the households have tap water with lead concentrations of more than $50 \mu\text{g litre}^{-1}$ (DHSS, 1980). Based on an average daily intake of $1.25 \text{ litre water day}^{-1}$, the daily intake of lead from water containing $50 \mu\text{g litre}^{-1}$ is $62 \mu\text{g}$.

Most of the lead in urban air originates from the combustion of lead-containing petrol. Organic lead in the form of tetraethyl- or tetramethyl lead is added to petrol in order to increase the octane number. The added amount of lead is regulated and, in most European countries, the maximum permitted concentration is $0.15 - 0.40 \text{ g litre}^{-1}$. Lead from car exhausts can be directly inhaled or deposited, or indirectly ingested by children through intake of soil or dust or through contaminated foods. Chamberlain et al. (1978) have studied the increase in blood lead level in relation to the environmental concentration of lead in air. They concluded that a concentration of lead in air of $1 \mu\text{g m}^{-3}$ will, through direct inhalation, increase the level of lead in blood by $2 \mu\text{g}/100 \text{ ml}$.

3.5 Contribution of lead from marine food

Relatively little data is available on lead concentrations in marine food. Table 11 gives a few examples of concentrations in fish and shellfish from the USA. Cod and flounder had levels of lead comparable to meat and poultry (about $0.1 \mu\text{g g}^{-1}$). Fresh salmon and mackerel had a higher content of lead. From the FAO/WHO Collaborating Centres for Food Contamination Monitoring (FAO/WHO, 1982), levels in fresh fish in the range of $0.04 - 0.3 \mu\text{g g}^{-1}$ were reported. Fresh crab and lobster from different

As for other kinds of food, concentrations in canned marine foods are much higher than in the corresponding fresh foods. Table 11 shows a more than 2-fold increase in lead concentrations in canned as opposed to fresh products.

3.6 Evaluation of potential health effects

The main source of lead intake for most individuals is through diet. A provisional tolerable weekly intake (PTWI) of 3.0 mg of lead in diet for adults was suggested by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) (FAO/WHO, 1972). This intake level is not applicable to children, and any increase in the amount of lead derived from drinking-water or inhaled from the atmosphere will reduce the amount tolerated through food intake. As PbB is the best indicator of total lead exposure and can be related to the health effects of lead, it would be of great importance to be able to relate lead intake to PbB levels. However, there are no conclusive data for such a relationship. Estimates of dietary lead absorption and retention can be applied; however, there are several modifying and interfering conditions. It is also likely that the relationship between ingested lead and PbB is not linear. The lack of data points to a great need for further studies in order to obtain information on the relationship between ingested lead and PbB.

PbB levels vary in different parts of the world, and mean levels are usually in the range of 10 - 20 $\mu\text{g}/100\text{ ml}$. Ninety percentile values above 25 $\mu\text{g}/100\text{ ml}$ have been reported in many countries.

The main target tissues for damage due to lead exposure are the haematopoietic and nervous systems. The major concern about low-level lead exposure is the risk of fetuses and children developing central nervous system dysfunction. These effects include those on behaviour, learning ability, intelligence, and fine motor co-ordination. Several steps in the haem biosynthetic pathway are affected by lead. Reduced activity of ALA-D, which has been observed at PbB levels above 10 $\mu\text{g}/100\text{ ml}$, is considered to be an indicator of elevated exposure rather than an adverse health effect. Increased levels of FEP have been connected with PbB levels of about 20 - 30 $\mu\text{g}/100\text{ ml}$ in adult females, about 25 - 35 $\mu\text{g}/100\text{ ml}$ in adult males, and about 15 $\mu\text{g}/100\text{ ml}$ in children. The dose levels which produce anaemia are not clearly demonstrated. In children, lead encephalopathy has occurred at PbB levels above 60 $\mu\text{g}/100\text{ ml}$. There are considerable uncertainties concerning the neuropsychological effects of low-level lead exposure in children due to methodological difficulties and because so many factors other than lead exposure can affect the intellectual development of children. The available data indicate that neuropsychological effects occur at PbB levels of more than 35 $\mu\text{g}/100\text{ ml}$, but more recent studies suggest harmful effects can occur at lower lead levels.

4. Conclusions on Lead

4.1 Potential harm to living resources

Lead inputs to the marine environment are derived from anthropogenic sources, including mining and the combustion of coal, wood, and other organic matter. The widespread use of alkyl-lead compounds in gasoline contributes to a diffuse atmospheric source which, following deposition or runoff, contributes in addition to the contamination of freshwater discharged to estuaries.

It has been estimated that as much as 90% of the atmospheric input to the sea may be of anthropogenic origin. There is evidence from analyses of inshore sediments that lead inputs have increased in recent years. However, levels of lead in surface offshore seawater is low, and recent analyses give concentrations of 10 ng litre^{-1} . Much of the lead input to the inshore waters appears to be in sediments. This lead, to an unknown extent, is likely to be mobilized during dredging and dredged spoil dumping.

Alkyl-lead compounds are readily degraded in sea water and are not likely to be persistent. Although there is some indication that alkyl lead compounds can be formed in the environment by microbial action, this has not been shown to be a very significant factor in lead recycling.

gammarid crustacea were affected at concentrations of 100 µg/litre. Reports on the higher sensitivity of 2 algal species need to be confirmed. Even so, available data do not show that the present levels of lead in the sea constitute a hazard to marine biota. Although there are no reported observations of effects of lead in marine mammals, it cannot be excluded that these may show biochemical effects from lead corresponding to those observed in terrestrial mammals.

Marine organisms, especially molluscs, can accumulate lead from contaminated environments, but the relative importance of water, sediments, and food as pathways for lead contamination is poorly understood. There is some evidence that birds may accumulate organic lead from contaminated molluscs, but the availability of such accumulated lead to other predators is unknown.

4.2 Potential hazards to human health

Dietary intake of lead, through food and drink, is the major contributor to the total body burden of lead in the general population. Other sources of exposure include water and air, and there is a possibility of exposure through lead in paints, dust, soil, and glazed domestic vessels. About 10% of the ingested lead in food is absorbed in the gastrointestinal tract of adults. Higher absorption, about 60%, has been shown under fasting conditions. In infants (up to 2 years of age), an absorption of about 40% has been reported. It is noted, however, that there is not only wide individual variability but several other factors which affect absorption.

Body burden of lead is divided into 2 fractions, the one firmly bound to bone and the other loosely bound to blood and soft tissues. The bone fraction, which constitutes about 90% of the total body burden of lead, has a biological half-time of approximately 5 years. The mean retention time of lead in blood and soft tissues is within the range of 3 weeks to 1 month.

The adult dietary intake of lead varies in different countries from about 30 µg day⁻¹ in Sweden to 400 µg day⁻¹ in Italy. There is a scarcity of data on the concentration of lead in fish. Average concentrations in the range of 0.1 - 0.4 mg kg⁻¹ fresh weight have been reported. In addition, some marine foods are canned for distribution, in which case lead from the cans may increase lead intake approximately 2-fold. It would normally be expected that fish does not greatly contribute to the daily intake of lead. The present PTWI dates as far back as 1972 and was based on a number of uncertain assumptions. The value is therefore considered to require re-evaluation, particularly as concerns pregnant women.

All contributions to environmental lead exposure are, thus, of concern. The neurotoxic effects of lead in human beings and, particularly, the developmental impairments in fetuses and children suggest that control of lead discharges to the environment, together with minimizing human exposure, are required.

5. References

ALEXANDER, F.W., DELVES, H.T., & CLAYTON, B.E. (1973) The uptake and excretion by children of lead and other contaminants. In: Proceedings of the International Symposium on the Environmental Health Aspects of Lead, Amsterdam, 2 - 6 October, 1972, Luxembourg, Commission of the European Communities, pp. 319-330.

ALEXANDER, F.W., CLAYTON, B.E., & DELVES, H.T. (1974) Mineral and trace-metal balances in children receiving normal and synthetic diets. Q. J. Med., 43: 89-111.

AMIARD, J.-C., AMIARD-TRIQUET, C., METAYER, C., MARCHAND, J., & FERRE, R. (1980) Etude du transfert de Cd, Pb, Cu, et Zn dans les chaines trophiques néritiques et estuariennes. I. Etat dans l'estuaire interne de la Loire (France) au cours de l'été 1978. Water Res., 14: 665-673.

AMIARD-TRIQUET, C., METAYER, C., AMIARD, J.-C., & FERRE, R. (1980) Etude du transfert de Cd, Pb, Cu, et Zn dans les chaines trophiques néritiques et estuariennes. II. Accumulation biologique chez les poissons planctonophages. Water Res., 14: 1327-1332.

- BAKER, E.L., LANDRIGAN, P.J., BARBOUR, A.G., COX, D.H., FOLLAND, D.S., LIGER, R.N., & THROCKMORTON, J. (1979) Occupational lead poisoning in the United States: clinical and biochemical findings related to blood lead levels. Br. J. ind. Med., 36: 314-322.
- BECKMAN, L., NORDENSON, I., & NORDSTROM, S. (1982) Occupational and environmental risks in and around a smelter in northern Sweden. VIII. Three-year follow-up of chromosomal aberrations in workers exposed to lead. Hereditas, 96: 261-264.
- BENIJTS-CLAUS, C. & BENIJTS, F. (1975) The effect of low lead and zinc concentration on the larval development of the mud crab Rhithropanopeus harrissi Gould. In: Koeman, J.H. & Strik, J.J.T.W.A., ed. Sublethal effects of toxic chemicals on aquatic animals, Amsterdam, Elsevier, pp. 43-52.
- BERLAND, B.R., BONIN, D.J., KAPKOV, V.I., MAESTRINI, S.Y., & ARLHAC, D.P. (1976) Action toxique de quatre métaux lourdes sur la croissance d'algues unicellulaires marines. C. R. Acad. Sci. Paris, 282: 633-636.
- BERLIN, A. (1982) Assessment of exposure to lead of the general population in the European Community through biological monitoring. Environ. Monit. Assess., 2: 225-231 (Presented at the Workshop on Integrated Exposure Assessment Monitoring, Las Vegas 19-23 Oct., 1981).
- BERNHARD, M. (1980) The relative importance of lead as a marine pollutant. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 345-352.
- BERTINE, K.K. (1980) Lead and the historical sedimentary record. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 319-324.
- BETTS, P.R., ASTLEY, R., & RAINE, D.N. (1973) Lead intoxication in children in Birmingham. Br. med. J., 1: 402-406.
- BIRKHEAD, R.B. (1982) Causes of mortality in the mute swan, Cygnus olor on the River Thames. J. Zool. Lond., 198: 15-25.
- BORNSCHEIN, R., PEARSON, D., & REITER, L. (1980) Behavioural effects of moderate lead exposure in children and animal models. Part I. Clinical studies. Part II. Animal studies. CRC crit. Rev. Toxicol., 43-99.
- BRANICA, M. & KONRAD, Z., ed. (1980) Lead in the marine environment, Oxford, Pergamon Press, 353 pp.
- BROWN, B. & AHSANULLAH, M. (1971) Effect of heavy metals on mortality and growth. Mar. Pollut. Bull., 2: 182-187.
- BRYAN, G.W. (1976) Some aspects of heavy metal tolerance in aquatic organisms. In: Lockwood, A.P.M., ed. Effects of pollutants on aquatic organisms, Cambridge, Cambridge University Press, pp. 7-34.
- BULL, K.R., EVERY, W.J., FREESTONE, P., HALL, J.R., & OSBORN, J. (1983) Alkyl lead pollution and bird mortalities on the Mersey estuary, UK, 1971-1981. Environ. Pollut. Ser. A., 31: 239-259.
- BURNETT, M. & PATTERSON, C.C. (1980) Analysis of natural and industrial lead in marine ecosystems. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 31-43.
- CALABRESE, A. & NELSON, D.A. (1974) Inhibition of embryonic development of the hard clam, Mercenaria mercenaria, by heavy metals. Bull. environ. Contam. Toxicol., 11: 92-97.
- CALABRESE, A., COLLIER, R.S., NELSON, D.A., & MCINNES, J.R. (1973) The toxicity of heavy metals to embryos of the American oyster Crassostrea virginica. Mar. Biol., 18: 162-166.

CAMBRAY, R.S., JEFFRIES, D.F., & TOPPING, G. (1975) An estimate of atmospheric trace elements inputs into the North Sea and the Clyde Sea (1972-1973), Harwell, Atomic Energy Authority, 30 pp (Report No. AERE, R7733).

CANTERFORD, G.S. & CANTERFORD, D.R. (1980) Toxicity of heavy metals to the marine diatom Ditylum brightwellii (West) Grunow: correlation between toxicity and metal speciation. J. Mar. Biol. Assoc. UK, 60: 227-242.

CHAISEMARTIN, C., CHAISEMARTIN, R., & BRETON, J. (1978) Aspect de la détresse métabolique chez macropodia: bioconcentration du plomb et activité de l'aspartate amino-transferase. C.R. Soc. Biol., 172: 1180-1187.

CHAMBERLAIN, A.C., HEARD, M.J., LITTLE, P., NEWTON, D., WELLS, A.C., & WIFFEN, R.D. (1978) Investigations into lead from motor vehicles, Harwell, Environmental and Medical Sciences Division (AERE - R 9198).

CHAU, Y.K. & WONG, P.T.S. (1980) Biotransformation and toxicity of lead in the aquatic environment. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 225-231.

CHISOLM, J.J., Jr & O'HARA, D.M. (1982) Lead absorption in children, Baltimore, Munich, Management, Clinical and Environmental Aspects. Urban and Schwarzenberg.

CHOW, T.J. (1978) Lead in natural waters. In: Nriagu, J.O., ed. The biogeochemistry of lead in the environment, Amsterdam, Elsevier, pp. 185-218.

CHOW, T.J. & PATTERSON, C.C. (1962) The occurrence and significance of lead isotopes in pelagic sediments. Geochim. Cosmochim. Acta, 26: 262-308.

CHOW, T.J., EARL, J.L., & BENNET, C.F. (1969) Lead aerosols in the marine atmosphere. Environ. Sci. Technol., 3: 737-740.

CHOW, T.J., EARL, J.L., & SNYDER, C.B. (1972) Lead aerosols in the marine atmosphere. Environ. Sci. Technol., 3: 737-740.

DAVIES, I.M. (1981) Survey of trace elements in fish and shellfish landed at Scottish Ports, 1975-76, 28 pp (DAFS Scottish Fisheries Research Report No. 19) (Department of Agriculture and Fisheries for Scotland).

DENTON, G.R.W. & BURDEN-JONES, C. (1981) Influence of temperature and salinity on the uptake, distribution, and depuration of mercury, cadmium, and lead by the Black-lip oyster, Saccostrea echinata. Mar. Biol., 64: 317-326.

DHSS (1980) Lead and health, London, Her Majesty's Stationery Office (The Report of a Department of Health and Social Security Working Party on Lead in the Environment).

DOE (1974) Report on lead in the environment, London, Department of the Environment, 4 pp (DOE Circular 115/74).

DUCE, R.A., HOFFMANN, G.L., FASCHING, J.R., & MOYERS, J.L. (1974) The collection and analysis of trace metals in atmospheric particulate matter over the North Atlantic. In: Proceedings of the WMO-WHO Technical Conference on the Observation and Measurement of Atmospheric Pollution, Helsinki, 30 July - 4 August 1973, Geneva, World Meteorological Organization, pp. 368, 370-379.

EGOROV, V.V., ZHIGALOVSKAJA, T.N., & MALAKHOV, S.G. (1970) Microelement content of surface air above the continent and the ocean. J. geophys. Res., 75(18): 3650-3656.

EISLER, R. (1983) Trace metal concentrations in marine algae, New York, Pergamon Press.

ELDON, J., PEKKARINEN, M., & KRISTOFFERSSON, R. (1980) Effects of low concentrations of heavy metals on the bivalve Macoma balthica. Ann. Zool. Fennici, 17: 233-242.

ERNHART, C.B., LANDA, B., & SCHELL, N.B. (1981) Subclinical levels of lead and development deficit-A multivariate follow-up reassessment. Pediatrics, 67: 911-919.

FAO/WHO (1972) Evaluation of certain food additives and the contaminants mercury, lead, and cadmium. Sixteenth Report of the Joint FAO/WHO Expert Committee on Food Additives, Geneva, World Health Organization (Technical Report Series No. 505).

FAO/WHO (1982) Summary and assessment of data received from the FAO/WHO Collaborating Centres for Food Contamination Monitoring, Uppsala, National Food Administration, pp. 61.

FLANAGAN, P.R., CHAMBERLAIN, M.J., & VALBERG, L.S. (1982) The relationship between iron and lead absorption in humans. Am. J. clin. Nutr., 36: 823-829.

FORSTNER, U. & VAN LIERDE, J.H. (1979) Trace metals in water purification process. In: Förstner, U. & Wittman, G.T.W., ed. Metal pollution in the aquatic environment, Berlin, Springer-Verlag, Chapter G., pp. 324-359.

FRIEDMAN, B. & HUTCHINSON, T.C. (1981) Sources of metal and elemental contamination of terrestrial environments. In: Lepp, N.W., ed. Effect of heavy metal pollution on plants, Barking, England, Pollution Monitoring Series, Applied Science, Vol. 2, pp. 35-94.

FUKAI, R. (1980) Input of lead into the Mediterranean through rivers. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 149-153.

GEORGE, S.G. (1980) Correlation of metal accumulation in mussels with the mechanisms of uptake, metabolism, and detoxification: a review. Thalassia. Jugosl., 16: 347-365.

GOLDBERG, E.D., PARKER, P.L., BOWEN, V.T., RISEBROUGH, R.W., FARRINGTON, J.W., ROBERTSON, W., HARVEY, G., SCHNEIDER, E., MARTIN, J.H., & GAMBLE, E. (1978) The mussel watch. Environ. Conserv., 5: 101-125.

GRAY, J.S. & VENTILLA, R.J. (1973) Growth rates of sediment-living marine protozoan as a toxicity indicator for heavy metals. Ambio, 2: 118-121.

HAMMOND, P.B., O'FLAHERTY, E.J., & GARTSIDE, P.S. (1981) The impact of air-lead on blood-lead in man: a critique of the recent literature. Food Cosmet. Toxicol., 19: 631-638.

HARRISON, R.M. & LAXEN, D.P.H. (1978) Natural source of tetraalkyllead in air. Nature, 275: 738-740.

HEARD, M.Y. & CHAMBERLAIN, A.C. (1982) Effects of minerals and food on uptake of lead from the gastrointestinal tract in humans. Human Toxicol., 1: 411-415.

HOLLIBAUGH, J.T., SEIBERT, D.L.R., & THOMAS, W.H. (1980) A comparison of the acute toxicities of ten heavy metals to phytoplankton from Saanich Inlet, B.C., Canada. Estuarine coastal Mar. Sci., 10: 93-105.

HRS-BRENKO, M., CLAUS, C., & BUBIC, S. (1977) Synergistic effects of lead, salinity, and temperature on embryonic development of the mussel, Mytilus galloprovincialis. Mar. Biol., 44: 109-115.

HUNTER, K.A. (1980) Processes affecting particulate matter in the sea surface microlayer. Mar. Chem., 9: 49-70.

IARC (1980) Some metals and metallic compounds, Lyons, International Agency for Research on Cancer, pp. 325-415 (Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans, Vol.23).

ICES (1974) Report of the Working Group for the International Study of the Pollution of the North Sea and its Effects on Living Resources and their Exploitation, Charlottenlund, Denmark, International Council for Exploration of the Sea (ICES Cooperative Research Report No. 39).

ICES (1977a) The ICES Coordinated Monitoring Programme in the North Sea, 1974, Charlottenlund, Denmark, International Council for Exploration of the Sea (ICES Cooperative Research Report No. 58).

ICES (1977b) A baseline study of the level of contaminating substances in living resources of the North Atlantic, Charlottenlund, Denmark, International Council for Exploration of the Sea (ICES Cooperative Research Report No. 69).

ICES (1977c) The ICES Coordinated Monitoring Programme, 1975 and 1976, Charlottenlund, Denmark, International Council for Exploration of the Sea (ICES Cooperative Research Report No. 72).

ICES (1980) Extention to the baseline study of contaminant levels in living resources of the North Atlantic, Charlottenlund, Denmark, International Council for Exploration of the Sea (ICES Cooperative Research Report No. 95).

IRPTC (1981) Data profile on lead in the marine and estuarine environment, Geneva, International Register of Potentially Toxic Chemicals (UNEP).

JAWORSKI, J.F. (1979) Effects of lead in the environment, 1978. Quantitative aspects, Canada, National Research Council (Publication No. NRCC 16736).

JELINEK, C.F. (1982) Levels of lead in the United States food supply. J. Assoc. Off. Anal. Chem., 65: 942-946.

JERNIGAN, E.L., RAY, B.J., & DUCE, R.A. (1971) Lead and bromine in atmospheric particulate matter on Oahu, Hawaii. Atmos. Environ., 5: 881-886.

JOHNSON, R.D., JONES, R.L., HINESLY, T.D., & DAVID, D.J. (1974) Selected chemical characteristics of soils, forages, and drainage water from the sewage farm serving Melbourne, Australia, Corps of Engineers, Department of the Army, USA, 54 pp.

JOHNSON, W.L. & DARMON, B.L. (1982) Influence of lead acetate or lead shot ingestion upon White Chinese Geese. Bull. enviromm. Contam. Toxicol., 29: 177-183.

JORHEM, L. & SLORACH, S.A. (1979) [Konservburkar av plat: källa för tenn och bly i livsmedel.] Tincans: a source of tin and lead in foodstuffs. Var Föda, 31: 173-191 (in Swedish with English summary).

KOBAYASHI, N. (1971) Fertilized sea urchin eggs as an indicatory material for marine pollution bioassay: preliminary experiments. Publs Seto mar. Biol. Lab., 18: 379- 406.

KOEPPE, D.E. (1981) Lead; understanding the minimal toxicity of lead in plants. In: Lepp, N.W., ed. Effects of heavy metal pollution on plants, Barking, England, Pollution Monitoring Series, Applied Science, Vol. 1., pp. 55-76.

KOSTIAL, K., SIMONOVIC, I., & PISONIC, U. (1971) Lead absorpction from the intestine in newborn rats. Nature (London), 233: 564.

KRAJNOVIC-OZRETIC, M. & OZRETIC, B. (1980) The ALA-D activity test in lead-exposed grey mullet, Mugil auratus. Mar. Ecol. Prog. Ser., 3: 187-191.

LANSDOWN, R., YULE, W., URBANOWICZ, M.-A., & MILLAR, I.B. (1983) Relationships between blood lead, intelligence, attainment and behaviour in school children: overview of a pilot study. In: Rutter, M. & Russel-Jones, R., ed. Lead and health, New York, John Wiley and Sons, Ltd., pp. 267-296.

LARSSON, B., SLORACH, S.A., HAGMAN, U., & HOFVANDER, Y. (1981) WHO collaborative breast feeding study. II. Levels of lead and cadmium in Swedish human milk, 1978-1979. Acta Paediatr. Scand., 70: 281-284.

LUOMA, S.N. & BRYAN, G.W. (1978) Factors controlling the availability of sediment bound lead to the estuarine bivalve, Scrobicula plana. J. Mar. Biol. Assoc. UK, 58: 793-802.

LUTZ, G.A., LEVIN, A.A., BLOOM, S.G., NIELSON, K.J., GROSS, J.L., & MORRISON, D.L. (1970) Lead model case study, Columbus, Ohio, Battelle Memorial Institution, Vol.3.

MADDOCK, B.G. & TAYLOR, D. (1980) The acute toxicity and bioaccumulation of some lead alkyl compounds in marine animals. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 233-261.

MAHAFFEY, K.R. (1981) Lead in the environment, Presented at the American Association for Advancement in Science Meeting, Toronto, Canada, 2-4 January 1981.

MAHAFFEY, K.R., ANNEST, J.L., ROBERTS, J., & MURPHY, R.S. (1982) National estimates of blood lead levels: United States, 1976-1980. Association with selected demographic and socioeconomic factors. N. Engl. J. Med., 307: 573-579.

MARCHETTI, R. (1978) Acute toxicity of alkyl leads to some marine organisms. Mar. Pollut. Bull., 9: 206-207.

MARSHALL, A.T. & TALBOT, V. (1979) Accumulation of cadmium and lead in the gills of Mytilus edulis: X-ray microanalysis and chemical analysis. Chem-Biol. Interact., 27: 111-123.

MARTIN, M., OSBORN, K.E., BILLIG, P., & GLICKSTEIN, N. (1981) Toxicities of ten metals to Crassostrea gigas and Mytilus edulis embryos and Cancer magister larvae. Mar. Pollut. Bull., 12: 305-308.

MILAR, C.R., SCHROEDER, S.R., MUSHAK, P., DOULCOURT, J.L., & GRANT, L.C. (1980) Contributions of the caregiving environment to increased lead burden of children. Am. J. Ment. Defic., 84: 339-344.

MOORE, M.R., MEREDITH, P.A., CAMPBELL, B.C., GOLDBERG, A., & PADDOCK, S.J. (1977) Contribution of lead in drinking-water to blood-lead. Lancet, 661-662.

MUROZUMI, M., CHOW, T.J., & PATTERSON, C.C. (1969) Chemical concentrations of pollutant lead aerosols, terrestrial dusts, and sea salts in Greenland and Antarctic snow strata. Geochim. Cosmochim. Acta, 33: 1247-1291.

MURRAY, A.J. (1979) Metals, organochlorine pesticide and PCB residue levels in fish and shellfish landed in England and Wales during 1974, Lowestoft, MAFF, 11 pp (Aquatic Environment Monitoring Report No. 2).

MURRAY, L.S., NORTON, M.G., NUNNY, R.S., & ROLFE, M.S. (1980) The field assessment of effects of dumping of wastes at sea. VI. The disposal of sewage sludge and industrial waste off the River Humber, Lowestoft, MAFF Fisheries, 35 pp (MAFF Research Technical Report No. 55).

NASH, W.W., POOR, B.W., & JENKINS, K.D. (1981) The uptake and subcellular distribution of lead in developing sea urching embryos. Comp. Biochem. Physiol., 69: 205-211.

NEEDLEMAN, H.L., ed. (1980) Low level lead exposure: the clinical implications of current research, New York, Raven Press.

NEEDLEMAN, H.L., GUNNOE, C., LEVITON, A., REED, R., PERESIE, H., MAHER, C., & BARRETT, P. (1979) Deficits in psychologic and classroom performance of children with elevated dentine lead levels. New Engl. J. Med., 300: 13.

NELMES, A.J., BUXTON, R. ST.J., FAIRWEATHER, F.A., & MARTIN, A.E. (1974) The implication of the transfer of trace metals from sewage sludge to man. In: Hemphill, D.D., ed. Proceedings of the 7th Annual Conference on Trace Substances and Environmental Health, Columbia, Missouri, University of Missouri, pp. 145-153.

NF (1982) Assessment of the safety of lead and lead salts in food, Nutrition Foundation (A Report of the Nutrition Foundation's Expert Advisory Committee).

NOIRFALISE, A. & COLLINGE, A. (1982) Metals and fluorides in wines. Toxicol. Eur. Res. 4: 201-204.

- NORDMAN, C.H. (1975) Environmental lead exposure in Finland. A study on selected population groups, Helsinki (Academic dissertation).
- NRIAGU, J.O., ed. (1978) The biogeochemistry of lead in the environment, Amsterdam, Elsevier (2 Vols., A & B).
- OLSEN, N.B., HOLLINAGEL, H., & GRANDJEAN, P. (1981) Indicators of lead exposure in an adult Danish suburban population. Dan. med. Bull., 28: 168-176.
- OSKARSSON, A. & CAMNER, P. (1983) Lead. In: Ewetz, L. & Camner, P., ed. Motor vehicles and cleaner air, Stockholm, Sweden, Liber Tryok, (Health risks from exposure to motor vehicle exhaust. A Report to the Swedish Governments Committee on Motors Air Pollution from the National Institute of Environmental Medicine).
- OTTO, D.A., BENIGNUS, V.A., MULLER, K.E., & BARTON, C.N. (1981) Effects of age and body lead burden on CNS function in young children. I. Slow cortical potentials. Electroencephalogr. clin. Neurophysiol., 52: 229-239.
- PATTERSON, C.C. (1965) Contaminated and natural lead environments of man. Arch. environ. Health, 11: 344-363.
- PATTERSON, C.C. & SEATTLE, D.M. (1976) The reduction of orders of magnitude errors in lead analysis of biological materials and natural waters by evaluating and controlling the extent and sources of industrial lead contamination introduced during sample collection and analysis. In: LaFleur, F.D., ed. Accuracy in trace analysis: sampling, sample handling, and analysis, Washington DC, US Department of Commerce, Vol. 1, pp. 321-334 (NBS Special Publication No. 422).
- PHILLIPS, D.J.H. (1976) The common mussel Mytilus edulis as an indicator of pollution by zinc, cadmium, lead, and copper. I. Effects of environmental variables on uptake of metals. Mar. Biol., 38: 59-69.
- PIOMELLI, S., CORASH, L., CORASH, M.B., SEAMAN, C., MUSHAK, P., GLOVER, B., & PADGETT, R. (1980) Blood lead concentrations in a remote Himalayan population. Science, 210: 1135-1137.
- PIOMELLI, S., SEAMAN, C., ZULLOW, D., CURRAN, A., & DAVIDO, W.B. (1982) Threshold for lead damage to heme synthesis in urban children. Proc. Natl Acad. Sci., 79: 3335- 3339.
- POOLE, C., SMYTHE, L.E., & ALPERS, M. (1980) Blood lead levels in Papua New Guinea children living in a remote area. Sci. Total Environ., 15: 17-24.
- RATCLIFFE, J.M. (1981) Lead in man and the environment, England, Ellis Horwood Ltd.
- RAY, S., MCLEESE, D.W., & PETERSON, M.R. (1981) Accumulation of copper, cadmium, and lead from two contaminated sediments by three marine invertebrates: a laboratory study. Bull. environ. Contam. Toxicol., 26: 315-322.
- REISH, D.J. & CARR, R.S. (1978) The effect of heavy metals on the survival, reproduction, development, and life cycles for two species of polychaetous annelids. Mar. Pollut. Bull., 9: 24-27.
- REISH, D.J., MARTIN, J.M., PILTZ, F.M., & WORD, J.W. (1976) The effect of heavy metals on laboratory populations of two polychaetes with comparisons to the water quality conditions and standards in Southern California marine waters. Water Res., 10: 299-302.
- RICE, D.C. (1982) Central nervous effects of perinatal exposure to lead or methylmercury in the monkey. Paper presented at the 15th Annual Rochester Conference, Rochester, New York, May 1982.
- RICKARD, D.T. & NRIAGU, J.O. (1978) Aqueous environmental chemistry of lead. In: Nriagu, J.O., ed. The biogeochemistry of lead in the environment, Amsterdam, Elsevier, pp. 219-284.

ROBINSON, I.M. (1978) Lead as a factor in the world economy. In: Nriagu, J.O., ed. The biogeochemistry of lead in the environment, Amsterdam, Elsevier, pp. 99-118.

ROELS, H.A., LAUWERYS, R.R., BUCHET, J.P., & VRELUST, M.-TH. (1975) Response of free erythrocyte prophyrin and urinary S-aminolevulinic acid in men and women moderately exposed to lead. Int. Arch. Arbeitsmed., 34: 97-108.

ROM, W.N. (1976) Effects of lead on the female and reproduction. A review. Mt. Sinai J. Med., 43: 542-552.

RUTTER, M. (1980) Raised lead levels and impaired cognitive/behavioural functioning: a review of the evidence. Dev. Med. Child. Neurol. (suppl No. 42): 1-26.

RUTTER, M. & RUSSEL-JONES, R. (1983) Lead versus health. Sources and effects of low-level exposure, New York, John Wiley and Sons, Ltd.

SCHAFFNER, R.M. (1981) Lead in canned foods. Food Technol., 35: 60-64.

SCHAULE, B. & PATTERSON, C.C. (1980) The occurrence of lead in the NE Pacific and the effects of ocean dumping on the New York Bight ecosystem. In: Ketchum, B.N., Kester, D.R., & Kihlo Park, P., ed. Ocean dumping of industrial wastes, New York, Plenum Press, pp. 53-86 (Marine Science Series 12).

SCHULZ-BALDES, M. (1974) Lead uptake from seawater and food, and lead loss in the common mussel, Mytilus edulis. Mar. Biol., 25: 177-195.

SCHUTZ, A. (1979) Cadmium and lead. Scand. J. Gastroenterol., 14(suppl. 52): 223-231.

SCHUTZ, A., SKERFVING, S., GULLBERG, B., & HAEGER-ARONSEN, B. (1981) Elimination of lead from blood after end of exposure. Paper presented at the XX International Congress on Occupational Health, Cairo, September 25- October 1, 1981.

SHAPER, A.G., POCOCK, S.J., WALKER, M., WALE, C.J., CLAYTON, B., DELVES, H.T., & HINKS, L. (1982) Effects of alcohol and smoking on blood lead in middle-aged British men. Br. med. J., 284: 299-302.

SHUSTER, C.N. & PRINGLE, B.H. (1969) Trace metal accumulation by the American Eastern Oyster, Crassostrea virginica. Proc. Natl Shell Fish. Assoc., 59: 91-103.

SILBERGELD, E.K. (1982) Neurochemical and ionic mechanisms of lead neurotoxicity. In: Prasad, K.N. & Vernadakis, A., ed. Mechanisms of actions of neurotoxic substances, New York, Raven Press.

SLORACH, S., GUSTAFSSON, I.-B, JORHEM, L., & MATTSSON, P. (1982) Intake of lead, cadmium, and certain other metals via a typical Swedish diet. Submitted for publication in Hum. Nutr. appl. Nutr., 1982.

SOMERO, G.N., YANCY, P.H., CHOW, T.S., & SNYDER, C.B. (1977) Lead effects on tissue and whole organism respiration of the estuarine teleost fish, Gillichthys mirabilis. Arch. environ. Contam. Toxicol., 6: 337-348.

STANFORD, H.M., O'CONNOR, J.S., & SWANSON, R.L. (1981) The effects of ocean dumping on the New York Bight ecosystem. In: Ketchum, B.H., Kester, D.R., & Kihlo Park, P., ed. Ocean dumping of wastes, New York, Plenum Press, pp. 53-86.

STEWART, J. & SCHULZ-BALDES, M. (1976) Long-term lead accumulation in abalone (Haliotis spp) fed on lead treated brown algae (Egrecia laevigata). Mar. Biol., 36: 19-24.

TASKINEN, H., NORDMAN, H., & HERNBERG, S. (1981) Blood lead levels in Finnish preschool children. Sci. total Environ., 20: 117-129.

TAYLOR, D. (1981) A summary of the data on the toxicity of various materials to

TOPPING, G., BEWERS, J.M., & JONES, P.G.W. (1980) A review of past and present measurements of selected trace metals in sea waters in the Oslo Commission and ICNAF/NAFO areas, 43 pp (ICES Cooperative Research Report No. 97).

TSUCHIYA, K. (1979) Lead. In: Friberg, L., Nordberg, G.F., & Vouk, V.B., ed. Handbook on the toxicology of metals, pp. 451-484, North Holland, Elsevier.

US EPA (1977) Air quality criteria for lead, Washington DC, US Environmental Protection Agency.

US EPA (1980) Ambient water quality criteria for lead, Washington DC, US Environmental Protection Agency, Criteria and Standards Division, 161 pp (US EPA No. 440/5-80-057).

VAHTER, M., ed. (1982) Assessment of human exposure to lead and cadmium through biological monitoring, Stockholm, Liber Tryck (Prepared for United Nations Environment Programme and World Health Organization by National Swedish Institute of Environmental Medicine and Department of Environmental Hygiene, The Karolinska Institute, Stockholm).

VARO, P. & KOIVISTOINEN, P. (1980) Mineral element composition of Finnish foods. XII. General discussion and nutritional evaluation. Acta Agric. Scand. (suppl. 22): 165-171.

WATERMAN, A.J. (1937) Effects of salts of heavy metals on development of the sea urchin, Arbacia punctulata. Biol. Bull., 73: 401-420.

WEIS, J.S. & WEIS, P. (1979) Effects of mercury, cadmium, and lead compounds on regeneration in estuarine fishes and crabs. In: Vernberg, W.B., Thurberg, F.R., Calabrese, A., & Vernberg, F.J., ed. Marine pollution: functional responses, New York Academic Press, pp. 151-169.

WHITFIELD, M., TURNER, D.R., & DICKSON, A.G. (1981) Speciation of dissolved constituents in estuaries. In: River inputs to ocean systems. Proceedings of UNESCO/IOC/UNEP Review Workshop, Rome, 16-19 March, 1979, New York, United Nations, pp. 132-188.

WHO (1977) Environmental health criteria 3: lead, Geneva, World Health Organization.

WHO (1980) Recommended health-based limits in occupational exposure to heavy metals, Geneva, World Health Organization (Report of a WHO Study Group) (Technical Report Series No. 647).

WINNEKE, G., KRAMER, U., BROCKHAUS, A., EWERS, U., KUJANEK, G., LECHNER, H., & JANKE, W. (1983) Neurophysiological studies in children with elevated tooth-lead concentrations. Int. Arch. occup. environ. Health, 51: 231-252.

WONG, P.T. (1975) Methylation of lead in the environment. Nature (London), 253: 263-264.

WOOD, J.M. (1980) Lead in the marine environment: some biochemical considerations. In: Branica, M. & Konrad, Z., ed. Lead in the marine environment, Oxford, Pergamon Press, pp. 299-303.

WOOLERY, M. & LEWIN, R.A. (1976) The effects of lead on algae. IV. Effects of Pb on respiration and photosynthesis of Phaeodactylum tricornutum (Bacillariophyceae). Water Air Soil Pollut., 6: 25-31.

YULE, W., LANDSDOWN, R., MILLAR, I.B., & URBANOWICZ, M.-A. (1981) The relationship between blood lead concentrations, intelligence and attainment in a school population: a pilot study. Dev. Med. Child. Neurol., 23: 567-576.

ZENCIRCI, N. (1980) Contribution à l'étude de l'accumulation et de la toxicité de l'étain et du plomb chez des crustacea gammarides. Hydrobiologia, 69: 179-186.

ZIEGLER, E.E., EDWARDS, B.B., JENSEN, R.L., MAHAFFEY, K.R., & FOMON, S.J. (1978) Absorption and retention of lead by infants. Pediatr. Res., 12: 29-34.

IV. TIN

1. Tin in the Marine Environment

1.1 Reference documentation

Apart from the individual papers referred to in the text and listed at the end, the major reference sources employed for this review were Environmental Health Criteria No. 15, Tin and Organotin Compounds (WHO, 1980), Handbook of Geochemistry (Wedepohl, 1969), and Trace Metal Concentrations in Marine Algae (Eisler, 1983).

1.2 General facts

Tin, with chemical symbol Sn (Latin Stannum), was discovered over 4000 years ago. The element belongs to Group IVb of the Periodic Table along with carbon, silicon, germanium, and lead. Its atomic number is 50, and its atomic weight is 118.70. Oxidation states in the environment are Sn^{2+} and Sn^{4+} . Recent studies (Pettine et al., 1981) show the predominant forms of Sn^{2+} in aerobic sea water at pH 8.1 to be $\text{Sn}(\text{OH})_2$, 93.8%, $\text{Sn}(\text{OH})_3$, 2.4%, and $\text{MeSn}(\text{OH})_3$, 3.8%. The final stable form in the sea may be as SnO_2 in bottom deposits. Sn^{2+} ionic forms must first be changed to hydrated oxides and then to Sn^{4+} hydrated oxides with a variable ionic configuration which is carried to the bottom. The oxidation-reduction potential of -0.13 V places redox reactions well within the physiological range. Tin can also be biomethylated in the environment, enhancing its toxicity and retention in biota (Hodge et al., 1979). Although ranking 21st in abundance among the 30 most familiar trace metals, tin is the eighth among trace metals in the human body. Schwartz (1974) has suggested that tin is an essential element for animals (rats), and there is evidence that it may be essential to man (Hamilton, 1979).

From considerations of the amount of input to the sea and its abundance in sea water and marine sediments, the nominal residence times quoted are 10^4 - 10^5 years (Förstner & van Lierde, 1979). More recently, Li et al. (1980) assessed the half-life of tin in the upper 350 m as only 3.5 years.

1.3 Sources

The main sources of tin are placer deposits in or derived from rivers, estuaries, and immediate offshore waters. The global distribution of commercially-useful deposits is shown by centres of production. World production in 1971 was distributed among Malaysia (32.3%), the USSR (12.1%), Bolivia (12.0%), Thailand (9.3%), China (8.7%), Indonesia (8.5%), Australia (4.1%), and others (13.0%) (which include Nigeria, Zaire, Brazil, and numerous minor contributors). Cassiterite (SnO_2) deposits, containing roughly 240 g tin m^{-3} , yield some 80% of the total production of 245 000 tonnes (1979) (Bauer, 1980); the remainder comes from various sulfide ores, some of which contain other metals as well as tin.

The mining, dredging, and beneficiation of tin are large scale operations requiring water, which inevitably causes the dispersal of tin-rich particulate matter in river and estuarine systems and, via them, to the sea. The cassiterite grains are 2.5 times more dense than sand and are separated by washing and sieving. Tin is recovered from ores by smelting techniques in which the tin is vaporized and condensed. As tin is very valuable, thorough recovery and reworking of tin residues and smelter fume is practiced.

In its metallic form, tin has many uses which promote virtually worldwide distribution; only the more massive and convenient products are retained for recycling. The main uses are as tinplate (much of which is unsuitable for re-use), solders and other alloys (bronze, babbitt-metal, pewter, type metal, dentists' amalgam, special air frame alloys), for strengthening glass, as a colour base, in catalysts, as a stabilizer for perfumes, and for sundry medical and dental applications (WHO, 1980; Peterson & Girling, 1981).

In recent years, synthetic tin compounds have increased to rank fourth among

pesticides, as catalysts, as antioxidants, in antifouling paints, and also to stabilize plastics and synthetic rubbers. Methyltins are considered biodegradable, yet complex synthetic organotins (e.g., triorganotin biocides) are resistant to environmental bacteria. Chemical attack requires either strongly acidic ($\text{pH} < 1$) or alkaline ($\text{pH} > 13$) conditions.

The passage of water through domestic plumbing and waste disposal systems greatly increases the level of tin in sewage. Trade waters from canning, dyeprinting, and laundries contain additional traces of tin.

The levels of tin (possibly including tin compounds) are relatively high in primary treated sewage, e.g., 4 - 6 mg litre⁻¹ in the United Kingdom and 110 - 170 $\mu\text{g g}^{-1}$ in dry weight sewage solids (Sterritt & Lester, 1980); 100 - 500 $\mu\text{g g}^{-1}$ in dry weight sewage solids is reported by Furr et al. (1976) for a series of USA samples. These values compare with only 3.8 $\mu\text{g g}^{-1}$ in cow manure.

Assuming a quantity of 150 $\mu\text{g g}^{-1}$ tin in dry weight sewage sludge (Förstner & von Lierde, 1979), the input of tin into the North Sea can be estimated at 1000 tonnes year⁻¹. A comparable amount could be assumed for the New York Bight (GESAMP, 1982). Clearly, even these approximate calculations demonstrate levels of local input that are important, although there is still no information about the contribution made by organotins. In the same way, large inputs of tin probably occur in channel and harbour dredgings which possess a high level of organic contamination.

1.4 Transport, transformation, and bioaccumulation

1.4.1 Transport

The rate of mobilization of tin by man ($240 \times 10^9 \text{ g year}^{-1}$) exceeds the rate of mobilization by natural forces 10-fold.

There is evidence that most of the transport of tin from the continents to the oceans is via the atmosphere. In the vicinity of tin smelters and recovery plants, airborne tin occurs at locally high concentrations (e.g., 3.8 - 4.4 $\mu\text{g m}^{-3}$ at 700 m from a Japanese smelter) (WHO, 1980) and some tens of ng m^{-3} , in polluted environments generally. The concentration in the marine air of the Northern Hemisphere is about 1 ng m^{-3} , but it is much less in the Southern Hemisphere (Byrd & Andreae, 1982).

Recent preliminary estimates of fluxes from the main atmospheric sources are given in Table 12 (Byrd & Andreae, 1982). Rates of natural and anthropogenic inputs into the global atmosphere appear to be substantial (12×10^9 to $70 \times 10^9 \text{ g year}^{-1}$), but estimates are speculative due to inadequate data. According to the authors, the high value attributed to waste incineration may be overestimated by up to a factor of 10.

Fluxes to the aquatic environment are just as uncertain; some examples are $0.22 \times 10^9 \text{ g year}^{-1}$ soluble tin in rivers and $83 \times 10^9 \text{ g yr}^{-1}$ tin attached to riverborne particles. This particulate tin is deposited almost totally in estuaries and inshore waters and contributes little to ocean input.

Freshwater transport of tin from land to sea originates from the weathering of low-level minerals, but the flux is reduced to a very low rate due to the characteristic, strong binding of tin and organotins to soil, sediment, and particulate matter. The average occurrence of tin in typical rocks is 0.5 $\mu\text{g g}^{-1}$ in sandstones, 2 $\mu\text{g g}^{-1}$ in igneous rocks, 3 - 4 $\mu\text{g g}^{-1}$ (range = 0.9 - 51 $\mu\text{g g}^{-1}$) in coals, and 6 $\mu\text{g g}^{-1}$ in shales. The concentration of tin in normal soils is usually between 1 and 10 $\mu\text{g g}^{-1}$ (Peterson & Girling, 1981).

1.4.2 Transformation

In recent years, many more analyses of soluble tin and organotins in freshwater have been reported. In most of the analyses of filtered waters reported by Byrd & Andreae (1982), mostly for southeastern USA rivers, total organotins average about 5% of total tin. An exception was the relatively high levels of organotin found in the German Rhine

Table 12. Estimated fluxes of tin into the atmosphere from natural and anthropogenic sources^a

Source	Global production or consumption (10 ¹² g/year)	Emission factor (g/g source)	Tin flux (10 ⁹ g/year)
<u>Anthropogenic input</u>			
Coal burning	3245	0.20 x 10 ⁻⁶	0.65
Coal burning	3245	0.17 x 10 ⁻⁶	0.55
Oil burning	1600	0.01 x 10 ⁻⁶	0.002
Wood and agricultural burning	1320	0.75 x 10 ⁻⁶ <u>b</u>	0.17
Wast incineration	540	8.60 x 10 ⁻⁵	47
Iron and steel production	1220	0.10 x 10 ⁻⁶ <u>c</u>	0.12
Non-ferrous metal production	18	3.50 x 10 ⁻⁴ <u>c</u>	6.3
Tin production	0.24	0.005 ^d (22)	1.2
Organotins	0.008	0.05	0.4
Estimated range			10 - 60
<u>Natural input</u>			
Sea spray	1000	2.80 x 10 ⁻¹⁰	0.0003
Soil dust	800	0.15 x 10 ⁻⁵	1.2
Volcanoes	25	0.96 x 10 ⁻⁵	0.24
Forest fires	320	0.75 x 10 ⁻⁶	0.24
Biomethylation			6 (?)
Estimated range			2 - 10
Total flux			12 - 70

^a From: Byrd & Andreae (1982).

^b Volatilization efficiency assumed to be 10%.

^c Emission efficiency assumed to be 10%.

^d Volatilization efficiency assumed to be 50%.

6 ng litre⁻¹ and 12 ng litre⁻¹, but the ranges were wide, 0.02 - 120 ng litre⁻¹ (Byrd & Andreae, 1982) and 1.3 - 37 ng litre⁻¹ with a single high value of 730 ng litre⁻¹ not included in the average (Braman & Tompkins). The single analysis of the polluted river Rhine (tin, 20 ng litre⁻¹; organotins, 300 ng litre⁻¹) shows a very high contribution from dimethyltin. Similarly, Hodge et al. (1979) (Table 13) report very high levels of tin in Lake Michigan with values of inorganic tin at 490 ng litre⁻¹ and 84 ng litre⁻¹, and butyltins at 2820 ng litre⁻¹ and 120 ng litre⁻¹ at 10 and 62 m depths, respectively.

In preliminary studies of the behaviour of soluble tin forms with respect to salinity in the estuary of the Ochlockonee River in Florida, Byrd & Andreae (1982) found some scavenging of inorganic tin and some indication that seawards, there were enhanced concentrations of methyltin compounds, especially dimethyltin. These data provide some field evidence for the formation of methyltins in estuaries. They highlight the marked local influence on tin and organotin levels of polluted freshwaters.

One of the earliest reports of biomethylation of tin comes from Huey et al. (1974), who observed the methylation of tin and of monomethyl tin by a Pseudomonas species. Subsequently, mono-, di-, and trimethyl tin compounds have been detected in freshwater, estuarine, and inshore waters (Braman & Tompkins, 1979 (Table 14); Hodge et al. 1979; Byrd & Andreae, 1982; Jackson et al., 1982), and biomethylation by certain bacteria in marine sediments has been repeatedly confirmed. Braman & Tompkins (1979) propose soil as a likely source of methyltins in freshwater.

The formation of inert volatile tetramethyltin from trimethyltin hydroxide is slow and not extensive (Guard et al., 1981), and it proceeds by biotic and abiotic action in marine sediments.

Bacterial demethylation has been demonstrated, and Jackson et al. (1982) suggest that the balance among methyltins in the environment represents a combination of methylation and demethylation activities. Exactly how aerobic and anaerobic conditions modify the formation of methyltins in sediments has not been fully explored.

1.4.3 Bioaccumulation

Very little information is available on the uptake of inorganic tin by marine organisms. Zencirci (1980) exposed the crustacean Gammarus locusta to 0.1 mg litre⁻¹ as stannic chloride and found that accumulation occurred on the cuticle and in the gut. However, as the solubility of tin in sea water is about 35 µg litre⁻¹, it is possible that this accumulation could have consisted of precipitated tin. Some information can be derived from data published by Smith & Burton (1972) who found that the southern United Kingdom coastal waters contained 0.02 - 0.04 µg litre⁻¹ of total tin. Algae (Phaeophyceae), gastropod, and lamellibranch molluscs from the same area contained 0.10 - 0.65, 0.33 - 0.71, and 0.23 - 0.67 µg tin g⁻¹ dry weight, respectively, indicating bioaccumulation factors within the range 2500 - 30 000 on a dry weight basis. Phytoplankton from Southampton Water contained 3.5 µg g⁻¹ dry weight, as compared to a concentration in the water of 0.04 µg litre⁻¹. In both cases, the contribution made by organotin compounds was unknown.

Dogan & Haerdi (1980) reported the concentrations of total tin in some plants, plankton, fishes, and sediments from one of the more contaminated Swiss freshwater lakes. By adopting a value of 500 ng litre⁻¹ for contaminated lake water (Hodge et al., 1979), the indicated biological accumulation factors for total tin would be in the range 6000 - 60 000 on a dry weight tissue basis. Representative values of the biological accumulation factors for methyltins alone are not yet available; however, the data of Dogan & Haerdi (1980) suggest a multiplier of less than 10 between plankton and fish tissue concentrations.

Few data are available for the bioaccumulation of organotin. Sheepshead minnows (Cyprinodon variegatus) exposed to 1.6 µg ¹⁴C-labelled tributyl tin oxide litre⁻¹ for 58 days did not reach an equilibrium within that period and, at the end, the bioaccumulation factor on a wet weight basis was 2600 for the whole fish (Ward et al., 1981). This was based on ¹⁴C measurements, and analysis showed that some breakdown to di-butyl and mono-butyl tin had occurred. There was a 78% loss of organotin after transfer to clean water for 28 days.

Table 13. Tin IV and organotin compounds in some environmental waters^a

Location	Collection date	Sn(IV)	MeSnCl ₃ - (ng/litre) ^b -	Me ₂ SnCl ₂
San Diego Bay, California (surface water)				
32°41'45"N 117°13'52"W	10/5/78	38 ± 2	8 ± 1	45 ± 2
32°43'05"N 117°13'32"W		6 ± 1	5 ± 1	38 ± 2
32°40'42"N 117°07'26"W ^c		9 ± 1	2 ± 1	15 ± 1
32°42'07"N 117°13'42"W		14 ± 1	2 ± 1	35 ± 1
32°40'57"N 117°13'42"W		13 ± 1	4 ± 1	20 ± 1
San Francisco Bay, California				
36°48'00"N 122°28'30"W (8 M)	10/14/78	2.1 ± 0.3	0	0
36°50'41"N 122°25'00"W (12 M)		3.2 ± 0.3	0	0
California coast, off San Francisco				
37°40'N 122°32'W (12 M)	10/11-12/78	0.8 ± 0.3	0	0
37°40'N 122°34'W (17 M)		0.5 ± 0.3	0	0
37°43'N 122°37'W (12 M)		0.3 ± 0.3	0	0

^a From: Hodge et al. (1979).

^b Organotin compounds are assumed to be chlorides for the calculation. 0 = not detected.

^c 10 ng/litre of a compound with the same retention time as the hydride of Et₂SnCl₂ found in the precipitate of this sample.

Table 14. Analysis of saline and estuarine water samples^{a,b}

Sample	Tin (IV) (ng litre ⁻¹) %		Methyl tin (ng litre ⁻¹) %		Dimethyl tin (ng litre ⁻¹) %		Trimethyl tin (ng litre ⁻¹) %		Total tin (ng litre ⁻¹)
<u>Saline waters</u>									
Gulf of Mexico, Sarasota	62.0	73	15	18	7.0	8.3	0.98	1.2	85
Gulf of Mexico, Fort Desoto	2.2	60	ND ^c		0.74	20	0.71	20	3.6
Gulf of Mexico, St Petersburg	4.5	54	0.62	7.4	3.2	39	ND	ND	8.3
Old Tampa Bay, Oldsmar	0.3	9.7	0.86	33	0.88	34	0.61	24	2.6
Old Tampa Bay, Safety Harbor	1.4	29	0.86	17	2.0	40	0.65	13	5.0
Old Tampa Bay, Philipee Park	0.8	32	1.1	44	0.60	24	ND	ND	2.5
Old Tampa Bay, Davis Municipal	ND	ND	0.98	35	0.91	32	0.95	34	2.8
Old Tampa Bay, Courtney Campbell	2.7	54	ND	ND	1.7	34	0.61	12	5.0
Average	1.7	40	0.63	15	1.4	33	0.50	12	4.2
<u>Estuarine surface waters</u>									
Sarasota Bay	5.7	47	3.3	27	2.0	16	1.1	9.1	12
Tampa Bay	3.3	27	8.0	66	0.79	6.5	ND	ND	12
McKay Bay	20	88	ND	ND	2.2	9.6	0.45	2.0	23
Hillsborough Bay	ND	ND	ND	ND	1.8	71	0.71	29	2.5
Hillsborough Bay, Seddon Channel; north	12	86	0.74	5.3	0.91	6.6	0.35	2.5	14
Hillsborough Bay, Seddon Channel; south	13	83	ND	ND	2.4	15	0.31	1.9	16
Manatee River	4.8	61	1.4	17	1.1	14	0.65	8.2	7.9
Alafia River	3.4	73	ND	ND	0.75	16	0.55	12	4.7
Palm River ^d	567	98	ND	ND	4.6	0.80	4.0	0.69	576
Bowes' Creek	8.6	42	8.5	42	3.3	16	ND	ND	20
Average	7.9	63	2.4	19	1.7	14	0.46	3.7	12

^a From: Braman & Tompkins (1979).^b Data are averages of duplicates.^c ND: less than 0.01 ng litre⁻¹ for methyl tin compounds and 0.3 ng litre⁻¹ for inorganic tin.^d This set of values was not used in computing the average.

Alzieu et al. (1982) also found that organotin compounds were readily taken up and lost by oysters held in tanks containing surfaces painted with anti-fouling compounds. When tissue levels reached $110 \mu\text{g tin g}^{-1}$ dry weight, a sequence of active gel secretion on the interior surfaces of the shell followed by calcium deposition occurred to give a chambered effect. Environmental evidence for this effect by organotins used in anti-fouling paints is given by Alzieu et al. (1980). Laboratory studies showed that bioaccumulation factors for Crassostrea gigas and Ostrea edulis exposed to $0.15 \mu\text{g tributyl tin oxide litre}^{-1}$ were 6000 and 1500, respectively, reaching a plateau after 10 days exposure. Only 50% of the accumulated organotin was depurated, within 10 days, in clean water (Waldock et al., 1983).

1.5 Tin and organotin concentrations in seawater, sediments, and marine biota

1.5.1 Sea water

An average concentration of tin in surface waters of the eastern Atlantic Ocean of 10 ng litre^{-1} has been reported by Smith & Burton (1972). More recently, Hodge et al. (1979) have shown a pronounced gradient of inorganic tin from 38 ng litre^{-1} for inshore California water to less than 1 ng litre^{-1} for offshore water.

In the Ochlockonee River estuary, inorganic tin changed from about 1 to 3 ng litre^{-1} at the head of the estuary to about $1 - 4.5 \text{ ng litre}^{-1}$ at the mouth with a hint of an overall increase in dimethyltin (Byrd & Andreae, 1982). The authors found near-surface values of inorganic tin between 1 and 3 ng litre^{-1} for 4 stations in the Gulf Stream with a pronounced decrease over the upper few hundred meters. At a depth of about 3000 m , concentrations were negligible but increased to about $0.6 \text{ ng litre}^{-1}$ below 4500 m . The likely source of the high near-surface values is atmospheric input; various interpretations of the slightly enhanced values in deep water might include benthic input or the presence of a separate deep water mass. Mono-, di-, and trimethyl tin distributions in the Ochlockonee estuary show consistent increasing concentrations seawards with dimethyltin as the dominant form (Byrd & Andreae, 1982). Values at the seaward limit were $0.3 - 1.2 \text{ ng litre}^{-1} \text{ Me}_2\text{Sn}^{2+}$, $0.4 \text{ ng litre}^{-1} \text{ Me}_3\text{Sn}^{2+}$, and $0.15 \text{ ng litre}^{-1} \text{ MeSn}^{2+}$.

Much higher values were found for San Diego Bay: $6 - 38 \text{ ng litre}^{-1}$ inorganic tin, $2 - 8 \text{ ng litre}^{-1} \text{ MeSnCl}_3$ and $15 - 45 \text{ ng litre}^{-1} \text{ Me}_2\text{SnCl}_2$ (Hodge et al., 1979); concentrations in samples farther offshore were below the limit of detection (MeSn , $0.5 \text{ ng litre}^{-1}$; Me_2Sn , $0.5 \text{ ng litre}^{-1}$). Examples of reported concentrations of tin in coastal and estuarine waters of the USA are given in Tables 13 and 14. Concentrations of about $0.15 \mu\text{g tributyltin litre}^{-1}$ were found in English east coast estuaries utilized by pleasure crafts, with higher concentrations occurring in marinas (Waldock & Thain, 1983).

1.5.2 Sediments

Even after excluding coastal sites of known tin occurrence (prospected or operational), values of total tin in inshore sediments are generally much higher than in deep-sea sediments. Hodge et al. (1979) claim to have demonstrated a well-marked and statistically-significant increased tin depositional rate over the past 50 years in the upper part of sediment cores from Narragansett Bay by a dating technique using ^{210}Pb . Tin concentrations in the upper $0 - 10 \text{ cm}$ lie in the range $20 - 14 \mu\text{g g}^{-1}$ (dry weight). These concentrations are much lower than those found by Dogan & Haerdi (1980) for Lake Lemana ($55 \mu\text{g g}^{-1}$ in sandy samples and $90 \mu\text{g g}^{-1}$ in sediments rich in humus).

Chester (1965) reported the following concentrations of tin in marine sediments: igneous rocks, $2 \mu\text{g g}^{-1}$; near-shore sediments, $21 \mu\text{g g}^{-1}$; manganese nodules, $300 \mu\text{g g}^{-1}$; and deep-sea argillaceous clays (Pacific Ocean), $20 \mu\text{g g}^{-1}$.

Smith & Burton (1972) found mean values for tin in ultramafic rocks to be $0.8 \mu\text{g g}^{-1}$; in basalts, $1.7 \mu\text{g g}^{-1}$; in silicic rocks, $2.5 \mu\text{g g}^{-1}$; in red clays, $3.4 \mu\text{g g}^{-1}$; in amphibolites, $1.2 \mu\text{g g}^{-1}$; and in ferromanganese deposits, $0.2 - 5.8 \mu\text{g g}^{-1}$. All of these values were found in estuarine, shelf, and Atlantic sediments.

Krauskopf (1956) suggested the possibility of co-precipitation of tin forms with CaCO_3 in deep water accompanied by bio-magnification in calcareous skeletons of unicellular organisms. Turekian (1965) has made a study of the distribution of tin concentration relative to the particle size of CaCO_3 ooze from the deep Pacific Ocean. Particles greater than about 50 μm had, on average, less than 10 $\mu\text{g g}^{-1}$ tin. The tin-rich fractions (44 - 67 $\mu\text{g g}^{-1}$) tend to fall between 0.7 and 10 μm . Braman & Tompkins (1979) report average forms of tin in a collection of sea shells: total Sn, 0.88 ng g^{-1} , MeSn, 0.24 ng g^{-1} , Me_2Sn , 0.051 ng g^{-1} , Me_3Sn , < 0.01 ng g^{-1} dry weight. The values for a white coral were: total Sn, 1.00 ng g^{-1} ; MeSn, 0.20 ng g^{-1} , Me_2Sn , 0.21 ng g^{-1} ; and Me_3Sn , < 0.01 ng g^{-1} dry weight.

1.5.3 Marine biota

Values for tin in the marine biota given in the review by Eisler (1983) are listed in Table 15. Apart from the values quoted for smoked-canned oysters, all the concentrations of total tin in edible material are moderate. The concentrations of tin and organotins in the marine food web have not been studied in detail. Samples from the Caribbean region, including fish, showed up to 8.8 ng g^{-1} organotin, but most of the samples contained only 0.25 - 1.0 ng g^{-1} total tin (Sherman & Carlson, 1980). Hodge et al. (1979) found the average total tin for 3 macroalgae (Pelagophycus, Macrocystis, Eisenia) to be 0.87 $\mu\text{g g}^{-1}$ dry weight. Many more observations are needed to provide a useful framework for an assessment of the roles of tin and organotins in the marine environment.

2. Effects on Marine Biota

2.1 Reference documentation

This section is based on information given in original publications as listed in the reference section. No general reviews were found in the literature.

2.2 Effect on marine biota

2.2.1 Inorganic tin

There are very few data on the toxicity of inorganic tin to marine organisms.

Algae

In a study of microorganisms in Chesapeake Bay, Hallas & Cooney (1981) found that the microbial flora were resistant to tin, but the exposure concentration used, 75 mg tin litre^{-1} , was well in excess of the reported solubility of about 35 $\mu\text{g tin litre}^{-1}$, so that the true exposure concentration is in doubt. Saboski (1977) found that exposure of the diatom Nitzschia liebethrutti to 178 $\mu\text{g tin litre}^{-1}$ for 14 days caused frustular abnormalities; the significance of this response is uncertain.

Crustacea

Exposure of Gammarus locusta to 0.1 mg tin litre^{-1} resulted in a 34% mortality in 16 days and 100% in 24 days (Zencirci, 1980).

Fish

No mortalities were observed among dab (Limanda limanda) exposed to a nominal 1 mg tin litre^{-1} for 4 days (G. Mance, personal communication), but precipitation and settlement of tin was observed with 500 $\mu\text{g tin litre}^{-1}$ remaining in suspension and 35 $\mu\text{g litre}^{-1}$ in solution.

These data are of very limited value in assessing the potential effects of elevated tin concentrations on marine organisms. However, precipitated tin in suspension does not appear to be highly toxic, and saturated solutions of tin in sea water may be toxic to crustacea only after prolonged exposure.

Table 15. Reported values for total tin in the marine biota and food preparations (ng g⁻¹)^a

Algae and macrophytes	0.5 - 101.0 dry weight	mainly 1971
Coelenterata		
<u>Beröe</u>	7.0 ash weight	1959
<u>Cyanea</u>	4.0 ash weight	1959
<u>Pleurobrachia</u>	20.0 ash weight	1962
Corals (34 spp)	< 5.0 dry weight	1971
Mollusca		
<u>Clione</u>	20.0 ash weight	1959
Mixed edible tissues	0.3 - 2.0 wet weight	1978
<u>Mytilus</u>	1.3 - 7.1 dry weight	1977
<u>Omnastrephes</u>	3.0 ash weight	1959
Smoked-canned oysters	25 - 30 wet weight	1950
Crustacea		
<u>Calanus</u>	< 1.0 ash weight	1959
<u>Centropages</u>	50 ash weight	1959
Copepods	70 ash weight	1962
Mixed edible tissues	0.6 - 2.0 wet weight	1978
Chaetognatha		
<u>Sagitta</u>	20 ash weight	1959
<u>Sagitta</u>	400 ash weight	1962
Tunicata		
<u>Salpa</u>	8 ash weight	1959
Teleosta		
Fish livers (26 spp)	0.2 - 0.4 wet weight	1978
Fish muscle (35 spp)	0.4 - 0.5 wet weight	1978
Fish muscle (75 spp)	0.5 - 0.6 wet weight	1978
<u>Morone</u> , liver	0.33 wet weight	1979
<u>Morone</u> , muscle	0.30 wet weight	1979

^a Source: Eisler (1983), which gives original references. The dates are given as a guide to the likely status of methodological development.

2.2.2 Organic tin

A number of organic tin compounds have been used in anti-fouling paint formulations, and these leach directly into sea water. Since the purpose of these paints is to kill sessile organisms which settle on the surface, the release of such substances from areas of high boat density, such as harbours and marinas, is a potential hazard to organisms in the surrounding area. The main compounds used are the trialkyl and triphenyl tins, and these form the basis of the available marine toxicity data.

(a) Alkyl tins

Bacteria

Crustacea

Wright & Roosenberg (1982) exposed Stage 1 zoeal larvae of the crab Uca pugnator to trimethyl tin over a range of salinities and temperatures. Significantly greater mortalities occurred in test solutions containing 20 $\mu\text{g litre}^{-1}$ trimethyl tin as compared to mortality in the control. Using larval stage of the shore crab (Hemigrapsus nudus) and lobster (Homarus americanus), Laughlin & French (1980) obtained the following data:

		Exposure concentration ($\mu\text{g litre}^{-1}$)	LT ₅₀ (days)
Shore crab	trimethyl tin oxide	50	6.6
	triethyl tin oxide	50	4.8
	tripropyl tin oxide	25	5.1
	tributyl tin oxide	25	6.2
Lobster	tributyl tin oxide	5	about 3
	tributyl tin oxide	1	no significant mortality

Mortalities increased at the time of shedding of the outer (or cuticular) layer. This was after 7 - 10 days of exposure for shore crab and after 3 - 5 days of exposure for lobster, which may explain the apparently greater sensitivity of the latter. Similar results were obtained with larval Carcinus maenas when it was exposed to tributyl tin oxide, for which the 96-h LC₅₀ was 10 $\mu\text{g litre}^{-1}$. Larval brown shrimp (Crangon crangon) were more sensitive with a 96-h LC₅₀ of 1.5 $\mu\text{g litre}^{-1}$, although adults were 10 times more resistant (Thain, 1983).

Tests on the harpacticoid Nitocra spinipes in brackish waters (salinity 7 ‰) gave 96-h LC₅₀s of 2 $\mu\text{g litre}^{-1}$ for tributyl tin as both the fluoride and the oxide (Linden et al., 1979).

Molluscs

The correlation between enhanced environmental concentrations of organic tin compounds and shell thickening and chambering in molluscs has been mentioned in section 4.1.4.3. Excessive shell thickening and reduction in growth occurred in juvenile Crassostrea gigas exposed to 0.15 $\mu\text{g tributyl tin oxide litre}^{-1}$ for 8 weeks. This effect was increased when additional suspended matter was present in the water (Waldock & Thain, 1983). Similar environmental concentrations in the vicinity of high pleasure-craft density produced the same effect on relaid oysters (Waldock & Thain, personal communication). His & Robert (1980) found that the eggs of the oyster C. gigas exposed to tributyl tin acetate did not develop at 5 $\mu\text{g litre}^{-1}$; "D" larvae suffered a 30% mortality, and the survivors showed no growth at this concentration. Thain (1983) found that the 48-h LC₅₀ for developing embryos of C. gigas and Ostrea edulis was 1.6 and 2.3 $\mu\text{g tributyl tin oxide litre}^{-1}$, respectively.

Juvenile clams (Rangia cuneata) exposed to tributyl tin acetate at 20 $\mu\text{g litre}^{-1}$ showed 100% mortality in 11 days, whereas adult clams were not killed by 90 $\mu\text{g litre}^{-1}$ in 17 days (Good et al., 1980).

Fish

Bleak (Alburnus alburnus), when exposed to tributyl tin fluoride in brackish water (salinity 7 ‰) had a 96-h LC₅₀ of 6 - 8 $\mu\text{g litre}^{-1}$. Its 96-h LC₅₀ was 15 $\mu\text{g litre}^{-1}$ when it was exposed to tributyl tin oxide (Linden et al., 1979). Using a mixture of fish species (Gambusia affinis, Peocilia latipinna, and Fundulus grandis) in estuarine water, Good et al. (1980) found that 100% mortality occurred within 11 days of exposure to 20 $\mu\text{g tributyl tin acetate litre}^{-1}$. Life cycle studies with the sheepshead minnow, Cyprinodon variegatus, showed that 1 $\mu\text{g tributyl}$

of $2.1 \mu\text{g litre}^{-1}$, whereas the corresponding concentrations for adult sole and the armed bullhead (Agonus cataphractus) were 36 and $16 \mu\text{g litre}^{-1}$, respectively.

(b) Aryl tins

Algae

Callow et al. (1979) found that the photosynthetic activity of zoospores of Enteromorpha intestinalis was reduced by 50% when it was exposed to triphenyl tin chloride (as tin) at $0.6 \mu\text{g litre}^{-1}$, whereas zoospores of Ulothrix flacca were similarly affected by $600 \mu\text{g litre}^{-1}$. Subsequently, Callow & Evans (1981) showed that the chlorophyll content of the diatom Achanthes subsessilis decreased during a 5-day exposure to concentrations of triphenyl tin acetate greater than $6 \mu\text{g litre}^{-1}$ (as tin). Mortalities occurred at $120 \mu\text{g litre}^{-1}$. There was some evidence of acclimatization to this organic tin during the exposure period.

Crustacea

Linden et al. (1979) found that the 96-h LC_{50} for the harpacticoid Nitocra spinipes when it was exposed to triphenyl tin fluoride in a brackish water (7 ‰) was $8 \mu\text{g litre}^{-1}$.

Fish

Further experiments by Linden et al. showed that the 96-h LC_{50} for bleak (Alburnus alburnus) when it was exposed to triphenyl tin fluoride was $40 \mu\text{g litre}^{-1}$.

3. Human Health Aspects

3.1 Reference documentation

This section is based primarily on original publications listed in the reference section. However, publications reviewing the health effects of inorganic tin and organotin compounds, such as the WHO (1980) Environmental Health Criteria document on the subject, and reviews by Barnes & Stoner (1959), Piscator (1979), and Bennett (1981) have also been used.

3.2 Toxicokinetic properties

The kinetics of tin, including gastrointestinal absorption, depend on its chemical form.

3.2.1 Inorganic tin

Inorganic tin is poorly absorbed from the gastrointestinal tract. Hiles (1974) found that less than 1% of the environmentally-dominant Sn (IV) form was absorbed in the rat after oral administration. In man, no differences can usually be found between oral intake and faecal excretion of tin (Schroeder et al., 1964; Calloway & McMullen, 1966). According to Johnson & Greger (1982), however, the daily supplementation of diet with 50 mg tin increased urinary tin excretion from $29 \mu\text{g day}^{-1}$ to $122 \mu\text{g day}^{-1}$.

All human tissues, with the exception of those of newborn babies, contain appreciable amounts of tin, and the highest concentration is found in the lung (Schroeder et al., 1964; Anspaugh et al., 1971). Hamilton et al. (1972, 1973) found 0.8 mg kg^{-1} tin in rib bones and lungs, 0.4 in liver, 0.2 in kidneys, and $0.005 \text{ mg litre}^{-1}$ in blood, mainly in red blood cells. The presence of tin in airborne particles, and a longer-term exposure explains the age-dependent increase in the lung concentration of tin (Schroeder et al., 1964). Bone is the major accumulation site for absorbed tin. Clearance from soft tissues is fast, but from bone it is slow (Furchner & Drake, 1976). After a fast elimination phase, approximately half of the absorbed dose is cleared both in man and experimental animals with a half-time of 400 days (Bennett, 1981).

Absorbed tin is mainly eliminated in urine. Perry & Perry (1959) and Meltzer et al. (1962) found mean values of 16.6 and $18.0 \mu\text{g litre}^{-1}$ total tin in the urine of

3.2.2 Organotin compounds

The kinetics of organotin compounds have not been studied in man and only to very limited extent in animals. Some quantitative data on gastrointestinal absorption (Bridges et al., 1967; Piscator, 1979), but mainly the differences between oral and intraperitoneal LD₅₀ values (Stoner et al., 1955; Kimbrough, 1976; WHO, 1980), indicate that absorption decreases with the number of carbon atoms in the organic radical. In the case of lower alkyltin compounds, gastrointestinal absorption also decreases with the number of valencies occupied by inorganic anions.

Many of the alkylated tin compounds are dealkylated in vivo (Cremer, 1957; Bridges et al., 1967) and, at least in the case of tributyltin, inorganic tin is the final product (Iwai et al., 1981). As in the case of absorption, excretion, too, depends on the degree of alkylation; an increase in the number of alkyl groups tends to shift excretion from urine to bile (Bridges et al., 1967).

The clearance half-time of trimethyltin is approximately 14 - 20 days in the rat (Brown et al., 1979), but as trimethyltin and triethyltin have exceptionally high affinities for rat haemoglobin (Rose & Aldridge, 1968), this clearance half-time cannot be extrapolated to other species, including man.

The urinary excretion of tin was increased above 100 µg litre⁻¹ by toxic occupational exposure to triphenyltin (Manzo et al., 1981). Braman & Tompkins (1979) found that 18% of the tin in urine was in methylated forms, but the mean total urinary tin concentration in the 11 samples was only 1 µg litre⁻¹, which is very much lower than the concentration found by other researchers.

3.3 Health effects

There are qualitative and quantitative differences in the toxicities of different tin compounds.

3.3.1 Inorganic tin

The oral toxicity of inorganic tin is low due to low absorption and rapid excretion. Outbreaks of nausea, vomiting, and diarrhoea have allegedly been caused by the consumption of orange-based drinks containing 0.43 g litre⁻¹ tin, although in studies with human volunteers, even 0.73 g litre⁻¹ tin caused no ill effects (Benoy et al., 1971). It has been suggested that the gastrointestinal effect of tin correlates with the concentration of tin in the drink and not with the total amount of tin ingested (Benoy et al., 1971).

Inorganic tin can interfere with the gastrointestinal absorption of essential elements, including calcium and zinc. This effect was apparent in human volunteers given 50 mg tin per day in their diet (Johnson et al., 1982). Very much higher doses inhibited bone calcification in experimental animals (Yamaguchi et al., 1981) and decreased the compressive strength of the femoral bone (Ogoshi et al., 1981).

3.3.2 Organotin compounds

The intrinsic toxicities of several lower members of the trialkyltin series and triphenyltin are of the same order (Stoner, 1966), but there are great variations between the different compounds in the manifestation of systemic toxicity. Their common localized effect is irritation. Lyle (1958) observed eye and skin irritation and sometimes skin burns in workers handling tributyltin, dibutyltin, and triphenyltin. Similar irritative effects, including nasal irritation, were reported by others (WHO, 1980).

The use of triphenyltin acetate as a fungicide spray resulted in headache, nausea, and vomiting in 2 sprayers (Manzo et al., 1981). One of them had 48 µg litre⁻¹ tin in blood and more than 100 µg litre⁻¹ tin in urine. Exposure to a mixture of triphenyltin acetate and manganese dithiocarbamate produced similar symptoms in 6 other cases. Three of them also developed liver damage (WHO, 1980). However, as no liver damage was seen in guinea-pigs treated with highly toxic doses of triphenyltin acetate (Stoner, 1966), the hepatotoxicity of this organotin seems questionable.

The main target of trimethyltin and triethyltin is the central nervous system, but within the central nervous system their effects are different. There are 3 reported cases of trimethyltin intoxication. One chemist exposed to trimethyltin developed hyperactivity, insomnia, and absent-mindedness; his recovery was complete (Brown et al., 1979). Two other chemists, who worked in a pilot plant and were exposed to dimethyltin and trimethyltin, suffered from memory defect, loss of vigilance, insomnia, anorexia, and disorientation. These symptoms were ignored until the persons developed mental confusion with general epileptic seizures. Both patients recovered completely after removal from exposure (Fortemps et al., 1978). In experimental animals, trimethyltin produces symmetrical neuronal damage in selected brain areas, mainly in the hippocampus but also in the neurocortex. Tremors, prostration, hyperactivity, aggression, and convulsions are the main toxic manifestations. In surviving animals, recovery can be complete (Brown et al., 1979). The marmoset, gerbil, and hamster are more sensitive than the rat because their haemoglobin does not bind trimethyltin. Based on experiments on the marmoset, Aldridge et al. (1981) predicted that the toxic dose of trimethyltin to man is 3.0 mg kg^{-1} or less.

The primary effect of human triethyltin intoxication is cerebral oedema (Barnes & Stoner, 1959) and the main clinical manifestations are headache, vomiting, disordered equilibrium, and coma (Alajouanine et al., 1958). Diethyltin capsules contaminated with triethyltin and used against boils and carbuncles caused 102 lethal and 108 non-lethal intoxications (Alajouanine et al., 1958). As the triethyltin content of capsules were not constant, the exact dose could not be calculated (Barnes & Stoner, 1959). Children seemed to be more sensitive than adults (WHO, 1980).

In experimental animals, the main pathological lesion is also cerebral oedema (Magee et al., 1957). The experiments of Stoner et al. (1955) indicate that, in the rabbit, the toxic oral dose is approximately $3.0 \text{ } \mu\text{g kg}^{-1}$ triethyltin chloride. For the rat, the minimal toxic triethyltin concentration in drinking-water is $5.0 \text{ mg litre}^{-1}$. This concentration produced brain oedema after 2 months exposure (Dieckman & Butler, 1971). Because of its conversion to triethyltin (Cremer, 1958), tetraethyltin produces the same effects in experimental animals as triethyltin (Barnes & Stoner, 1959).

There are no reported cases of systemic intoxication caused by the higher alkyl analogues. On a molar basis, they are less toxic than the lower alkyl homologues and they do not produce hippocampal damage or brain oedema (Mushak et al., 1982). The same is true for dibutyltin. In experimental animals, dibutyltin salts are able to damage the biliary tract, liver, and pancreas (Barnes & Magee, 1958) and to decrease thymus weight. This latter effect is independent of the length of the alkyl radical (Henninghausen et al., 1980).

3.4 Total exposure to tin

The daily intake of tin by man ranges from 200 to 17 000 mg per head. As the contributions of water ($30 \text{ } \mu\text{g day}^{-1}$) and atmospheric pollution (less than $1 \text{ } \mu\text{g day}^{-1}$) are small, the major source of tin is food. Fresh food can provide 4.0 mg day^{-1} but, when food is stored in cans or PVC containers, significantly more tin can be ingested (WHO, 1980). Organotin pesticide residues in fresh foods have only a negligible effect on exposure (WHO, 1980).

3.5 Contribution of tin from marine food

There are few data on tin concentrations in fresh marine food, and these are not representative. The flesh of fish or molluscs caught in polluted bays and coastal areas may contain 1 mg kg^{-1} tin, and even higher concentrations can be present in fish from harbour areas (UNEP, 1981). Since per capita average daily fish consumption in Europe is 38 g day^{-1} (Hamilton and Minski, 1972/73), it seems reasonable to use 40 g day^{-1} fish consumption as a starting point for calculation. Fish consumption from products containing 1 mg kg^{-1} inorganic tin adds $40 \text{ } \mu\text{g}$ to the daily tin intake from fish. This contribution is small so that a 10-fold or even 100-fold increase in fish consumption could not result in toxic inorganic tin exposure. The increased consumption of marine food products will decrease the consumption of other types of foods and, consequently, tin intake from other sources. However, tin in fish and in other marine food products can be organic, both from anthropogenic sources and synthesized in marine

In harbours, bays, and off-shore areas, anthropogenic organotin biocides or their metabolites (e.g., monobutyltin from tri- and dibutyltin) can be present in appreciable concentration and can enter the bioaccumulation process (Hodge et al., 1979; Seidel et al., 1980; Jackson et al., 1982). A more ubiquitous source of organotin is the gradual methylation of inorganic tin to mono-, di-, tri-, and tetramethyltin (Braman & Tompkins, 1979; Byrd & Andreae, 1982; Jackson et al., 1982). The presence of methylated tin compounds in seashells (Braman & Tompkins, 1979), macro algae, and invertebrates (Seidel et al., 1980) has been demonstrated, and this makes their presence in marine food products probable. At present, lack of data prevents the assessment of any hazard to human health.

When marine food products are used for feeding livestock, the restricted absorption of tin from the gastrointestinal tract limits, as far as inorganic tin is concerned, the transfer of marine tin to meat products.

3.6 Evaluation of potential health effects

It seems very unlikely that inorganic tin in marine food products, directly or indirectly, presents any hazard to human health. However, neither exposures nor health hazard presented by the possible accumulation of organotin compounds, especially trimethyltin and tetramethyltin, can be assessed.

4. Conclusions on Tin

4.1 Potential harm to living resources

World production of tin is about 0.25 million tonnes per annum, and the substance is used widely both as metal and as organic tin compounds. Evidence from tin concentrations in estuarine sediments indicate that the discharge of tin to the marine environment has increased steadily during the past 100 years. Much of the input is derived from diffuse sources, such as domestic and industrial wastes (including those from tin-mining), which may be discharged via rivers or directly, including by the dumping of sewage sludge.

Specific point sources include offshore tin mining of those sediments in which primary tin deposits are present in a potentially soluble form (e.g., stannite and varlamoffite) as occurs in Malaysia.

The use of organic tin compounds in anti-fouling paints for ships may give rise to elevated concentrations in harbours and estuaries where shipping is concentrated.

Most of the data for tin concentrations in seawater refer to total tin and reported concentrations range from 1 to 5 ng litre⁻¹ in offshore waters with up to 40 ng litre⁻¹ in inshore waters. More recently, some analyses have been made of organotin concentrations, which are important because of the high toxicity of some of the different tin species and compounds.

Soluble inorganic tin entering estuaries may be adsorbed or co-precipitated on particles, and there is some evidence that such tin can become methylated by bacterial action. The extent to which this occurs is not known. Similarly, organotin compounds can be slowly degraded by abiotic and biotic processes but, again, there is little quantitative information on degradation rates.

The data available for this review indicates that inorganic tin and a range of organotin compounds have widely different toxic properties. Few experiments have been carried out with inorganic tin and these have been at concentrations above the solubility limit of 35 µg litre⁻¹. Neither precipitated tin nor saturated solutions have been shown to be acutely toxic to a limited range of marine organisms, although effects on crustaceans have been recorded with chronic exposure. Since this solubility limit is a 1000 times greater than recorded total tin concentrations inshore, it is unlikely that existing levels of contamination are approaching harmful concentrations.

Organotins used in anti-fouling paints are much more toxic than inorganic tin. Although the data on alkyl and aryl tins are not extensive, they are reasonably

range of species. Early life stages of organisms, particularly molluscs and moulting crustacea, may be harmed at lower concentrations.

The use of organotins in antifouling paints is a direct source of hazard to marine organisms in the vicinity of harbours, marinas, and shipping lanes. At present, only commercially-exploited shellfish are known to have been harmed, but it is likely that other species are also at risk. On the evidence available, it is not possible to derive a concentration below which the biota would not be harmed.

Depending on the extent to which methylation of inorganic tin occurs, areas containing tin-contaminated sediments, derived from domestic and industrial inputs, may form a potential source of toxic organotin compounds which could also occur in offshore mining areas.

4.2 Potential hazards to human health

Tin is acquired by many individuals through ingestion of food without occupational exposure. Tin in the food can be in inorganic or organic forms. The gastrointestinal absorption of inorganic tin is low, probably below 5%, and animal experiments indicate that the gastrointestinal absorption of the organotins with longer alkylchains is similarly low. There is only indirect evidence to support the suggestion that gastrointestinal absorption of the lower homologues is higher and increases with increasing alkylation.

In man, the highest concentrations of tin are found in bones and in lungs ($0.8 \mu\text{g g}^{-1}$), while concentrations in blood are less than one hundredth of this (5 ng ml^{-1}). Increases in oral intake of inorganic tin is not usually, but may be, associated with corresponding increases in faecal tin excretion and with some increases in urinary excretion.

At present, there is no established relationship between exposure to tin and the blood concentration or excretion rates. Depending on its chemical form, tin exerts different toxic effects. Acute inorganic tin intoxication after oral ingestion is manifested by nausea, vomiting, and diarrhoea caused by the local effect of tin on the gastrointestinal mucosa. A diet supplemented with 50 mg tin per day for 20 days had no adverse effect but did cause an increase in faecal zinc excretion.

The toxicity of organotin compounds shows wide variation. Cases of intoxication have been reported only for triethyltin, trimethyltin, and triphenyltin. The most toxic, trimethyltin and triethyltin, damaged the central nervous system, while triphenyltin caused only general symptoms such as headache, nausea, and vomiting. Children seem to be more sensitive to triethyltin than adults, but in none of the cases of organotin poisonings was dose defined or biological parameters monitored.

The dietary intake of tin shows wide variation depending on the regional distribution of tin in soil, the use of tin in industry, and its use in agriculture. The greatest increase in dietary tin, however, is caused by the release of tin from cans or plastic containers. The dietary intake of tin is increased by the predominant use of tinned food from less than 1 mg day^{-1} to well over 100 mg day^{-1} , but without any noticeable adverse effect.

The contribution of fresh fish or other seafood to the daily intake of tin is low. If one assumes 40 g day^{-1} fish consumption from products containing 1 mg kg^{-1} inorganic tin, the consumption of tin is only $40 \mu\text{g day}^{-1}$, a small proportion of the normal daily tin intake. However, tin in fish and in other marine food products can be organic, both from anthropogenic sources and synthesized by marine biota. The decomposition of anthropogenic organotin compounds in the marine environment, the synthesis of trimethyltin in the marine biota, and the irreversibility of the damage inflicted by doses as low as 3.0 mg kg^{-1} trimethyltin on the CNS in experimental animals, make trimethyltin the most important organotin compound in the marine environment in terms of potential toxicity. However, at present it is difficult to predict whether or not trimethyltin in marine food alone, or in conjunction with other tin compounds, exhibits any hazard to human health. The reason for this is 2-fold: (1) the lack of concentrated data on marine food products; and (2) the lack of quantitative human toxicological data both on toxic doses and clearance half-time.

5. References

ALAJOUANINE, T., DEROBERT, L., & THIERFY, S. (1958) Etude clinique d'ensemble de 210 cas d'intoxication par les sels organique d'étain. Rev. Neurol., 98: 85-96.

ALDRIDGE, W.N., BROWN, A.W., BRIERLEY, J.B., VERSCHOYLE, R.D., & STREET, B.W. (1981) Brain damage due to trimethyltin compounds. Lancet, II: 692-693.

ALZIEU, C., THIBAUD, Y., HERAL, M., & BOUTIER, B. (1980) Evaluation des risques dus à l'emploi des peintures anti-salissures dans les zones conchylicoles. Rev. Trav. Inst. Pêches Maritimes, 44: 305-348.

ALZIEU, C., HERAL, M., THIBAUD, Y., DARDIGNAC, M.-J., & FEUILLETT, M. (1982) Influence des peintures anti-salissures à base organostanniques sur la calcification de la coquille de l'huitre Crassostrea gigas. Rev. Trav. Inst. Pêches Maritimes, 45: (in press).

ANSPAUGH, L.R., ROBINSON, W.C., MARTIN, W.H., & LOWE, O. (1971) Compilation of published information on elemental concentrations in human organs in both normal and diseased states, University of California, Vol. 2, Pt. 2 (UCRL 51013).

BARNES, J.M. & MAGEE, P.N. (1958) The biliary hepatic lesion produced experimentally by dibutyltin salts. J. Pathol. Bacteriol., 75: 267-279.

BARNES, J.M. & STONER, H.B. (1959) The toxicology of tin compounds. Pharmacol. Rev., 11: 211-231.

BAUER, W., ed. (1980) Metal statistics 1969-1979, Frankfurt, Metallgesellschaft AG.

BENNETT, B.G. (1981) Exposure commitment assessments of environmental pollutants. Summary exposure assessments for mercury, nickel, tin, London, Chelsea College (MARC Report No.25).

BENOY, C.J., HOOPER, P.A., & SCHNEIDER, R. (1971) The toxicity of tin in canned fruit juices and solid foods. Food Cosmet. Toxicol., 9: 645-656.

BRAMAN, R.S. & TOMPKINS, M.A. (1979) Separation and determination of nanogram amount of inorganic tin and methyltin compounds in the environment. Anal. Chem., 51: 12-19.

BRIDGES, J.W., DAVIES, D.S., & WILLIAMS, R.T. (1967) The fate of ethyltin and diethyltin derivatives in the rat. Biochem. J., 105: 1261-1266.

BROWN, A.W., ALDRIDGE, W.N., STREET, B.W., & VERSCHOYLE, R.D. (1979) The behavioural and neuropathologic sequelae of intoxication by trimethyltin compounds in the rat. Am. J. Pathol., 97: 59-82.

BYRD, J.T. & ANDREAE, M.O. (1982) Tin and methyltin species in seawater: concentrations and fluxes. Science, 218: 565-569.

CALLOW, M.E. & EVANS, L.V. (1981) Some effects of triphenyl tin chloride on Achnanthes subsessilis. Bot. Mar., 24: 201-205.

CALLOW, M.E., MILLER, P.A., & EVANS, L.V. (1979) Organotin resistance in green seaweeds. In: Proceedings of the International Seaweed Symposium, Vol. 9, pp. 191-197.

CALLOWAY, D.H. & MCMULLEN, J.J. (1966) Fecal excretion of iron and tin by men fed stored canned foods. Am. J. clin. Nutr., 18: 1-6.

CHESTER, R. (1965) Geochemical criteria for the differentiation of reef and non-reef facies in carbonate rocks. Bull. Am. Assoc. Petrol. Geol., 49: 258-276.

CREMER, J.E. (1957) The metabolism in vitro of tissue slices from rats given triethyltin compounds. Biochem. J., 67: 87-96.

CREMER, J.E. (1958) The biochemistry of organotin compounds: the conversion of

DIECKMAN, W. & BUTLER, W.H. (1971) Histopathological studies and chronic toxic effects. In: Proceedings of the 12th Meeting of the European Society for the Study of Drug Toxicity, Vol. 12, pp. 24-27.

DOGAN, S. & HAERDI, W. (1980) Determination of total tin in environmental biological and water samples by AAS with graphite furnace. Int. J. environ. anal. Chem., 8: 249-257.

EISLER, R. (1983) Trace metal concentrations in marine algae, London, Pergamon Press.

FORSTNER, & VAN LIERDE, J.H. (1979) Trace metals in water purification proceses. In: Förstner, U. & Wittman, G.T.W., ed. Metal pollution in the aquatic environment, Berlin, Springer-Verlag, Chapter G., pp. 324-366.

FORTEMPS, E., AMAND, G., BOMBOIR, A., LAUWREYS, R., & LATERRE, E.C. (1978) Trimethyltin poisoning. Report of two cases. Int. Arch. occup. Health, 41: 1-6.

FURCHNER, J.E. & DRAKE, G.A. (1976) Comparative metabolism of radionuclides in mammals. XI. Retention of ^{113}Sn in the mouse, rat, monkey, and dog. Health Phys., 31: 219-224.

FURR, A.K., LAWRENCE, A.W., TONG, S.S.C., GRANDOLFO, M.C., HOFSTADER, R.A., BACHE, C.A., GUTENMANN, W.H., & LISK, D.J. (1976) Multi-element and chlorinated hydrocarbon analysis of municipal sludges of American cities. Environ. Sci. Technol., 10: 683-687.

GESAMP (1982) The health of the oceans, Geneva, United Nations Environment Programme, 108 pp (Rep. Stud. GESAMP No. 15).

GOOD, M.L., DUNDEE, D.S., & SWINDLER, G. (1980) Bioassay and environmental effects of organotin marine antifoulants. In: Baker, R., ed. Controlled release of bioactive materials, Academic Press, pp. 387-397.

GUARD, H.E., COBET, A.B., & COLEMAN, W.M. (1981) Methylation of trimethyltin compounds by estuarine sediments. Science, 213: 770-771.

HALLAS, L.E. & COONEY, J.J. (1981) Tin and tin-resident micro-organisms in Chesapeake Bay. Appl. environ. Microbiol., 41: 466-471.

HAMILTON, E.I. (1979) The chemical elements and man: measurements, perspectives, applications, Springfield, Illinois, C.C. Thomas, 528 pp.

HAMILTON, E.I. & MINSKI, M.J. (1972/73) Abundance of the chemical elements in man's diet and possible relations with environmental factors. Sci. total Environ., 1: 375-394.

HAMILTON, E.I., MINSKI, M.J., & CLEARY, J.J. (1972/73) The concentration and distribution of some stable elements in healthy human tissues from the United Kingdom. An environmental study. Sci. total Environ., 1: 341-374.

HENNIGHAUSEN, G., LANGE, P., & MERKORD, J. (1980) The relationship between the length of the alkyl chain of dialkyltin compounds and their effects on thymus and bile ducts in mice. Arch. Toxicol. (suppl. 4): 175-178.

HILES, R.A. (1974) Absorption, distribution, and excretion of inorganic tin in rats. Toxicol. appl. Pharmacol., 27: 366-379.

HIS, E. & ROBERT, R. (1980) Action d'un sel organometallique, l'acetate de tributyl-etain sur les oeufs et larves D de Crassostrea gigas (Thurnberg), Copenhagen, International Council for Exploration of the Sea (ICES CM 1980/F:27 (mimeo)).

HODGE, V.F., SEIDEL, S.L., & GOLDBERG, E.D. (1979) Determination of tin (IV) and organotin compounds in natural waters, coastal sediments, and macro algae by atomic absorption spectrometry. Anal. Chem., 51: 1256-1259.

HUEY, C., BRINKMAN, F.E., GRIM, S., & IVERSON, W.P. (1974) The role of bacterial

IWAI, H., WADA, O., & ARAKAWA, Y. (1981) Determination of tri-, di-, and monobutyltin and inorganic tin in biological materials and some aspects of their metabolism in rats. J. anal. Toxicol., 5: 300-305.

JACKSON, J.A.A., BLAIR, W.R., BRINCKMAN, F.E., & IVERSON, W.P. (1982) Gas-chromatographic speciation of methylstannates in Chesapeake Bay. Environ. Sci. Technol., 16: 110-119.

JOHNSON, M.A. & GREGER, J.L. (1982) Effects of dietary tin on tin and calcium metabolism of adult males. Am. J. clin. Nutr., 35: 655-660.

JOHNSON, M.A., BAIER, M.J., & GREGER, J.L. (1982) Effects of dietary tin on zinc, copper, iron, manganese, and magnesium metabolism in adult males. Am. J. clin. Nutr., 35: 1332-1338.

KIMBROUGH, R.D. (1976) Toxicity and health effects of selected organotin compounds: a review. Environ. Health Perspect., 14: 51-56.

KRAUSKOPF, K.B. (1956) Factors controlling the concentration of thirteen rare metals in sea water. Geochim. Cosmochim. Acta, 9: 1-32.

LAUGHLIN, R.B. & FRENCH, W.J. (1980) Comparative study of the acute toxicity of a homologous series of trialkyl tins to larval shore crabs, Hemigrapsus nudus, and lobster Homarus americanus. Bull. environ. Contam. Toxicol., 25: 802-809.

LI, Y.-H., FEELY, H.W., & TOGGWEILER, J.R. (1980) ^{228}Ra and ^{228}Th concentrations in GEOSECS Atlantic surface waters. Deep-Sea Res., 27: 545-555.

LINDEN, E., BENGTSSON, B.-E., SVANBERG, O., & SUNDSTROM, G. (1979) The acute toxicity of 78 chemicals and pesticide formulations against two brackish water organisms, the bleak (Alburnus alburnus) and the harpacticoid Nitocra spinipes. Chemosphere, 8: 843-851.

LYLE, W.H. (1958) Lesions of the skin in process workers caused by contact with butyltin compounds. Br. J. ind. Med., 15: 193-196.

MAGEE, P.N., STONER, H.B., & BARNES, J.M. (1957) The experimental production of oedema in the central nervous system of the rat by triethyltin compounds. J. Pathol. Bacteriol., 73: 107-124.

MANZO, L., RICHELMI, P., SABBIONI, E., PIETRA, R., BONO, F., & GUARDIA, L. (1981) Poisoning by triphenyltin. Report of two cases and determination of tin in blood and urine by neutron activation analysis. Clin. Toxicol., 18: 1343-1353.

MELTZER, L.E., RUTMAN, J., GEORGE, P., RUTMAN, R., & MITCHELL, J.R. (1962) Urinary excretion pattern of trace metals in diabetes mellitus. Am. J. med. Sci., 244: 282-289.

MUSHAK, P., KRIGMAN, M.R., & MALLIMAN, R.B. (1982) Comparative organotin toxicity in the developing rat: somatic and morphological relationship to accumulate total tin. Neurobehav. Toxicol., 4: 209-215.

OGOSHI, K., KURUMATANI, N., AOKI, Y., MORIYAMA, T., & NANZAI, Y. (1981) Decrease in compressive strength of the femoral bone in rats administered stannous chloride for a short period. Toxicol. appl. Pharmacol., 58: 331-332.

PERRY, H.M., Jr & PERRY, E.F. (1959) Normal concentrations of some trace metals in human urine: changes produced by ethylene-diaminetetraacetate. J. clin. Invest., 38: 1452-1463.

PETERSON, P.J. & GIRLING, C.A. (1981) Other trace metals: 12, Tin. In: Lepp, N.W., ed. Effect of heavy metal pollution on plants, London, Applied Science, Vol. 1, pp. 249-252 (Pollution Monitoring Series).

- PISCATOR, M. (1979) Tin. In: Friberg, L., Nordberg, G.F., & Vouk, V.B., ed. Handbook on the toxicology of metals, Amsterdam, Elsevier/North Holland Biomedical Press, pp. 613-626.
- ROSE, M.S. & ALDRIDGE, W.N. (1968) The interaction of triethyltin with components of animal tissues. Biochem. J., 106: 821-828.
- SABOSKI, E.M. (1977) Effects of mercury and tin on frustular ultrastructure of the marine diatom Nitzschia lieberthruetti. Water Air Soil Pollut., 8: 461-466.
- SCHROEDER, H.A., BALASSA, J.J., & TIPTON, I.H. (1964) Abnormal trace metals in man: tin. J. chron. Dis., 17: 483-502.
- SCHWARTZ, K. (1974) Recent dietary trace element research, exemplified by Sn, F, and Si. Fed. Proc., 33: 1748-1757.
- SEIDEL, S.L., HODGE, V.F., & GOLDBERG, E.D. (1980) Tin as environmental pollutant. Thalassia Yugosl., 16: 209-223.
- SHERMAN, L.R. & CARLSON, T.L. (1980) A modified phenylfluorone method for determining organotin compounds in ppb and sub-ppb ranges. J. anal. Toxicol., 4: 31-33.
- SMITH, J.D. & BURTON, J.D. (1972) The occurrence and distribution of tin with particular reference to marine environments. Geochim. Cosmochim. Acta, 36: 621-633.
- STERRITT, R.M. & LESTER, J.N. (1980) Determination of Ag, Co, Mn, and Sn in sewage sludge by a rapid electrothermal AAS method. Analyst (London), 105: 616-620.
- STONER, H.B. (1966) Toxicity of triphenyltin. Br. J. ind. Med., 23: 222-229.
- STONER, H.B., BARNES, J.M., & DUFF, J.I. (1955) Studies on the toxicity of alkyl tin compounds. Br. J. Pharmacol., 10: 16-25.
- THAIN, J.E. (1983) The acute toxicity of bis (tributyl tin) oxide to the adults and larvae of some marine organisms, International Council for Exploration of the Sea (ICES No. CM 1983/E:13).
- TUREKIAN, K.K. (1965) Some aspects of the geochemistry of marine sediments. I. In: Riley, J.P. & Skirrow, G., ed. Chemical oceanography, London, Academic Press, Vol. 2, Chapter 16.
- UNEP (1981) Data profile on tin in the marine and estuary environment, Geneva, United Nations Environment Programme (UNEP FP/0503-77-03).
- WALDOCK, M.J. & THAIN, J.E. (1983) Shell thickening in Crassostrea gigas: organotin antifouling or sediment induced? Mar. Pollut. Bull., 14: 411-415.
- WALDOCK, M.J., THAIN, J.E., & MILLER, D. (1983) The accumulation and depuration of bis (tributyl tin) oxide in oysters: a comparison between the Pacific oyster (Crassostrea gigas) and the European flat oyster (Ostrea edulis), International Council for Exploration of the Sea (ICES CM 1983/E:52).
- WARD, G.S., CRAMM, G.C., PARRISH, P.R., TRACHMAN, H., & SLESINGER, A. (1981) Bioaccumulation and chronic toxicity of bis (tributyl tin) oxide (TBTO): tests with a salt water fish. In: Branson, D.R. & Dickson, K.L., ed. Aquatic toxicology and hazard assessment: fourth conference, pp. 183-200 (ASTM STP 737).
- WEDEPOHL, K.H. (1969) Handbook of geochemistry, Berlin, Springer.
- WHO (1980) Environmental Health Criteria 15: tin and organotin compounds, Geneva, World Health Organization, pp. 109.
- WRIGHT, D.A. & ROSENBERG, W.H. (1982) Trimethyl tin toxicity to larval Uca pugilator: effects of temperature and salinity. Arch. environ. Contam. Toxicol., 11: 491-495.

YAMAGUCHI, M., SUGII, K., & OKADA, S. (1981) Change in the mineral composition and its related enzyme activity in the femur of rats orally administered stannous chloride. J. Pharmacol. Dynamics, 4: 874-878.

ZENCIRCI, N. (1980) Contribution à l'étude de l'accumulation et de la toxicité de l'étain et du plomb chez des crustacés gammarides. Hydrobiologia, 69: 179-186.

ANNEX I: FIRST SESSION OF THE GESAMP WORKING GROUP ON REVIEW OF POTENTIALLY HARMFUL SUBSTANCES: CADMIUM, LEAD, AND TIN (Geneva, 31 January - 4 February, 1983)

Participants

Dr B.G. Bennett, Monitoring and Assessment Research Centre, The Octagon Building,
London, United Kingdom

Prof L. Friberg, National Institute of Environmental Medicine, Department of Environment
Hygiene of the Karolinska Institute, Stockholm, Sweden (Chairman)

Prof A. Furtado Rahde, Brazilian Drug and Poison Information System, Fundaçao Oswaldo
Cruz, Ministry of Health, Porto Alegre, Brazil

Prof A. Jernelöv, Swedish Environmental Research Institute, Stockholm, Sweden

Mr R. Lloyd, Ministry of Agriculture, Fisheries, and Food, Fisheries Laboratory,
Burnham-on-Crouch, Essex, United Kingdom

Dr L. Magos, Medical Research Council, Toxicology Unit, Carshalton, Surrey, United
Kingdom

Prof. S.P. Meyers, Department of Food Science, Louisiana State University, Baton Rouge,
Louisiana, USA

Dr A. Oskarsson, National Institute of Environmental Medicine, Department of
Environmental Hygiene of the Karolinska Institute, Stockholm, Sweden

Dr P.M. Sivalingam, School of Biological Sciences, University of Sciences Malaysia,
Minden, Penang, Malaysia

Prof. G. Tomassi, National Institute of Nutrition, Rome, Italy

WHO Secretariat

Dr H. Galal-Gorchev, Environmental Hazards and Food Protection, World Health
Organization, Geneva, Switzerland

Dr M. Gilbert, International Register of Potentially Toxic Chemicals, United Nations
Environment Programme, Geneva, Switzerland

Dr R. Helmer, WHO Technical Secretary of GESAMP, World Health Organization, Geneva,
Switzerland

Dr J. Parizek, International Programme on Chemical Safety, World Health Organization,
Geneva, Switzerland

Dr G. Vettorazzi, International Programme on Chemical Safety, World Health Organization,
Geneva, Switzerland

Reports and Studies GESAMP

The following reports and studies have been published so far. They are available from any of the organizations sponsoring GESAMP.

1. Report of the seventh session, London, 24-30 April 1975. (1975). Rep.Stud.GESAMP, (1):pag.var. Available also in French, Spanish, and Russian.
2. Review of harmful substances. (1976). Rep.Stud.GESAMP, (2):80 p.
3. Scientific criteria for the selection of sites for dumping of wastes into the sea. (1975). Rep.Stud.GESAMP, (3):21 p. Available also in French, Spanish, and Russian.
4. Report of the eighth session, Rome, 21-27 April 1976. (1976). Rep.Stud.GESAMP, (4):pag.var. Available also in French and Russian.
5. Principles for developing coastal water quality criteria. (1976). Rep.Stud.GESAMP, (5):23 p.
6. Impact of oil on the marine environment. (1977). Rep.Stud.GESAMP, (6):250 p.
7. Scientific aspects of pollution arising from the exploration and exploitation of the sea-bed. (1977). Rep.Stud.GESAMP, (7):37 p.
8. Report of the ninth session, New York, 7-11 March 1977. (1977). Rep.Stud.GESAMP, (8):33 p. Available also in French and Russian.
9. Report of the tenth session, Paris, 29 May - 2 June 1978. (1978). Rep.Stud.GESAMP, (9):pag.var. Available also in French, Spanish, and Russian.
10. Report of the eleventh session, Dubrovnik, 25-29 February 1980. (1980). Rep.Stud.GESAMP, (10):pag.var. Available also in French and Spanish.
11. Marine Pollution implications of coastal area development. (1980). Rep.Stud.GESAMP, (11):114 p.
12. Monitoring biological variables related to marine pollution. (1980). Rep.Stud.GESAMP, (12):22 p. Available also in Russian.
13. Interchange of pollutants between the atmosphere and the oceans. (1980). Rep.Stud.GESAMP, (13):55 p.
14. Report of the twelfth session, Geneva, 22-29 October 1981. (1981). Rep.Stud.GESAMP, (14):pag.var. Available also in French, Spanish, and Russian.
15. The review of the health of the oceans. (1982). Rep.Stud.GESAMP, (15):108 p.
16. Scientific criteria for the selection of waste disposal sites at sea. (1982). Rep.Stud.GESAMP, (16):60 p.
17. The evaluation of the hazards of harmful substances carried by ships. (1982). Rep.Stud.GESAMP, (17):pag.var.
18. Report of the thirteenth session, Geneva, 28 February - 4 March 1983. (1983). Rep.Stud.GESAMP, (18):50 p. Available also in French, Spanish, and Russian.
19. An oceanographic model for the dispersion of wastes disposed of in the deep sea. (1983). Rep.Stud.GESAMP, (19):182 p.
20. Marine pollution implications of ocean energy development. (in press). Rep.Stud.GESAMP, (20).

22. Review of potentially harmful substances - Cadmium, lead and tin. (1985). Rep.Stud.GESAMP, (22):114 p.
23. Interchange of pollutants between the atmosphere and the oceans. (in press). Rep.Stud.GESAMP, (23).
24. Thermal discharges in the marine environment. (1984). Rep.Stud.GESAMP, (24).44 p.

PUBLICATIONS IN THE UNEP REGIONAL SEAS REPORTS AND STUDIES SERIES

- No. 1 UNEP: Achievements and planned development of UNEP's Regional Seas Programme and comparable programmes sponsored by other bodies. (1982)
- No. 2 UNIDO/UNEP: Survey of marine pollutants from industrial sources in the West and Central African region. (1982)
- No. 3 UNESCO/UNEP: River inputs to the West and Central African marine environment. (1982)
- No. 4 IMCO/UNEP: The status of oil pollution and oil pollution control in the West and Central African region. (1982)
- No. 5 IAEA/UNEP: Survey of tar, oil, chlorinated hydrocarbons and trace metal pollution in coastal waters of the Sultanate of Oman. (1982)
- No. 6 UN/UNESCO/UNEP: Marine and coastal area development in the East African region. (1982)
- No. 7 UNIDO/UNEP: Industrial sources of marine and coastal pollution in the East African region. (1982)
- No. 8 FAO/UNEP: Marine pollution in the East African region. (1982)
- No. 9 WHO/UNEP: Public health problems in the coastal zone of the East African region. (1982)
- No. 10 IMO/UNEP: Oil pollution control in the East African region. (1982)
- No. 11 IUCN/UNEP: Conservation of coastal and marine ecosystems and living resources of the East African region. (1982)
- No. 12 UNEP: Environmental problems of the East African region. (1982)
- No. 13 UNEP: Pollution and the marine environment in the Indian Ocean. (1982)
- No. 14 UNEP/CEPAL: Development and environment in the Wider Caribbean region: A Synthesis. (1982)
- No. 15 UNEP: Guidelines and principles for the preparation and implementation of comprehensive action plans for the protection and development of marine and coastal areas of regional seas. (1982)
- No. 16 GESAMP: The health of the oceans. (1982)
- No. 17 UNEP: Regional Seas Programme: Legislative authority. (in preparation)
- No. 18 UNEP: Regional Seas Programme: Workplan. (1982)
- No. 19 Rev. 2. UNEP: UNEP Oceans Programme: Compendium of projects. (1985)
- No. 20 CPPS/UNEP: Action Plan for the protection of the marine environment and coastal areas of the South-East Pacific. (1983)

- No. 21 CPPS/UNEP: Sources, levels and effects of marine pollution in the South-East Pacific. (1983) (In Spanish only)
- No. 22 Rev. 2. UNEP: Regional Seas Programme in Latin America and Wider Caribbean. (1985)
- No. 23 FAO/UNESCO/IOC/WHO/WMO/IAEA/UNEP: Co-ordinated Mediterranean Pollution Monitoring and Research Programme (MED POL) - Phase I: Programme Description. (1983)
- No. 24 UNEP: Action Plan for the protection and development of the marine and coastal areas of the East Asian region. (1983)
- No. 25 UNEP: Marine pollution. (1983)
- No. 26 UNEP: Action Plan for the Caribbean environment programme. (1983)
- No. 27 UNEP: Action Plan for the protection and development of the marine environment and coastal areas of the West and Central African region. (1983)
- No. 28 UNEP: Long-term programme for pollution monitoring and research in the Mediterranean (MED POL) - Phase II. (1983)
- No. 29 SPC/SPEC/ESCAP/UNEP: Action Plan for managing the natural resources and environment of the South Pacific region. (1983)
- No. 30 UNDIESA/UNEP: Ocean energy potential of the West and Central African region. (1983)
- No. 31 A. L. DAHL and I. L. BAUMGART: The state of the environment in the South Pacific. (1983)
- No. 32 UNEP/ECE/UNIDO/FAO/UNESCO/WHO/IAEA: Pollutants from land-based sources in the Mediterranean. (1984)
- No. 33 UNDIESA/UNEP: Onshore impact of offshore oil and natural gas development in the West and Central African region. (1984)
- No. 34 UNEP: Action Plan for the protection of the Mediterranean. (1984)
- No. 35 UNEP: Action Plan for the protection of the marine environment and the coastal areas of Bahrain, Iran, Iraq, Kuwait, Oman, Qatar, Saudi Arabia and the United Arab Emirates. (1983)
- No. 36 UNEP/ECLAC: The state of marine pollution in the Wider Caribbean region. (1984)
- No. 37 UNDIESA/UNEP: Environmental management problems in resource utilization and survey of resources in the West and Central African region. (1984)
- No. 38 FAO/UNEP: Legal aspects of protecting and managing the marine and coastal environment of the East African region. (1983)
- No. 39 IUCN/UNEP: Marine and coastal conservation in the East African region. (1984)

- No. 40 SPC/SPEC/ESCAP/UNEP: Radioactivity in the South Pacific. (1984)
- No. 41 UNEP: Socio-economic activities that may have an impact on the marine and coastal environment of the East African region. (1984)
- No. 42 GESAMP: Principles for developing coastal water quality criteria. (1984)
- No. 43 CPPS/UNEP: Contingency plan to combat oil pollution in the South-East Pacific in cases of emergency. (1984)
- No. 44 IMO/ROPME/UNEP: Combating oil pollution in the Kuwait Action Plan region. (1984)
- No. 45 GESAMP: Thermal discharges in the marine environment. (1984)
- No. 46 UNEP: The marine and coastal environment of the West and Central African region and its state of pollution. (1984)
- No. 47 UNEP: Prospects for global ocean pollution monitoring. (1984)
- No. 48 SPC/SPEC/ESCAP/UNEP: Hazardous waste storage and disposal in the South Pacific. (1984)
- No. 48/Appendices SPC/SPEC/ESCAP/UNEP: Hazardous waste storage and disposal in the South Pacific. (1984)
- No. 49 FAO/UNEP: Legal aspects of protecting and managing the marine and coastal environment of the East African region: National Reports. (1984)
- No. 50 IUCN/UNEP: Marine and coastal conservation in the East African region: National Reports. (1984)
- No. 51 UNEP: Socio-economic activities that may have an impact on the marine and coastal environment of the East African region: National Reports. (1984)
- No. 52 UNEP: Arab co-operation for the protection and development of the marine environment and coastal areas resources of the Mediterranean. (1984)
- No. 53 UNEP: UNEP Regional Seas Programme: the Eastern African Experience. (1984)
- No. 54 UNIDO/UNEP: Contingency planning for emergencies associated with industrial installations in the West and Central African region. (1985)
- No. 55 FAO/UNEP: Marine mammals: global plan of action. (1985)
- No. 55/Annex FAO/IUCN/IWC/UNEP: Marine mammals: global plan of action. (1985)
- No. 56 GESAMP: Cadmium, lead and tin in the marine environment. (1985)
- No. 57 IMO/UNEP: Oil spills and shoreline clean-up on the coasts of the Eastern African region. (1985)
- No. 58 UNEP: Co-operative programmes sponsored by UNEP for the protection of the marine and coastal environment in the wider Indian Ocean region. (1985)

- No. 59 UNEP: Environmental problems of the marine and coastal area of India: National Report. (1985)
- No. 60 IUCN/UNEP: Management and conservation of renewable marine resources in the Indian Ocean region: Overview. (1985)
- No. 61 UNEP: Action Plan for the protection, management and development of the marine and coastal environment of the Eastern African region. (1985)
- No. 62 IUCN/UNEP: Management and conservation of renewable marine resources in the South Asian Seas region. (1985)
- No. 63 IUCN/UNEP: Management and conservation of renewable marine resources in the Kuwait Action Plan region. (1985)
- No. 64 IUCN/UNEP: Management and conservation of renewable marine resources in the Red Sea region. (1985)
- No. 65 IUCN/UNEP: Management and conservation of renewable marine resources in the East Asian Seas region. (1985)
- No. 66 IUCN/UNEP: Management and conservation of renewable marine resources in the Eastern African region. (1985)
- No. 67 UN/UNEP: Coastal erosion in West and Central Africa. (1985)